



Mandel, R., Whay, H. R., Klement, E., & Nicol, C. J. (2016).  
Environmental enrichment of dairy cows and calves in indoor housing.  
*Journal of Dairy Science*, 99(3), 1695-1715.  
<https://doi.org/10.3168/jds.2015-9875>

Peer reviewed version

License (if available):  
Unspecified

Link to published version (if available):  
[10.3168/jds.2015-9875](https://doi.org/10.3168/jds.2015-9875)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1 INTERPRETIVE SUMMARY

2

3 Nearly all dairy cattle are housed indoors, at least for some part of their life. Keeping cows  
4 indoors allows farmers to meet the nutritional needs of high yielding individuals and provide  
5 protection from predators. However, it also confronts cows and calves with a wide range of  
6 environmental challenges such as restricted movement and exposure to loud aversive sounds.  
7 Here we review recent evidence of the effect of environmental enrichment on the welfare of  
8 dairy cattle kept indoors. This is accompanied by an assessment of the practicality of different  
9 enrichment options, considering the divergent needs of calves and cows separately.

**Environmental enrichment of dairy cows and calves in indoor housing**

**R. Mandel\*<sup>1</sup>, H.R. Whay†, E. Klement\*, and C.J Nicol†**

\*The Hebrew University, Jerusalem, Israel

†University of Bristol, Bristol, England

<sup>1</sup> Corresponding author: Roi Mandel, Koret School of Veterinary Medicine, Robert H. Smith

Faculty of Agriculture, Food and Environment, The Hebrew University, P.O.B. 12, Rehovot

76100, Israel. Phone (office) - 972-(0)8-9489560, Fax - 972-(0)8-9489138

**roi.mandel@mail.huji.ac.il**

## ABSTRACT

In recent years, there has been an increase in the number of farmers who choose to keep their cows indoors throughout the year. Indoor housing of cows allows farmers to provide high yielding individuals with a nutritionally balanced diet fit for their needs, and it has important welfare benefits for both cows and their calves, such as protection from predators, parasites and exposure to extreme weather conditions. However, it also confronts cows and calves with a wide range of environmental challenges. These include abiotic environmental sources of stress (e.g. exposure to loud and aversive sound) and confinement-specific stressors (e.g. restricted movement and maintenance in abnormal social groups). Cows and calves that live indoors are also faced with the challenge of occupying long periods of time with a limited range of possible behavioural patterns. Environmental enrichment can improve biological functioning (measured as increased lifetime reproductive success, increased inclusive fitness or a correlate of these such as improved health), help animals to cope with stressors in their surroundings, prevent frustration, increase the fulfillment of behavioural needs, and promote more positive affective states. Here we review recent findings on the effect of social, occupational, physical, sensory and nutritional enrichment on dairy cows and calves, and we assess the appropriateness and practicality of implementing different enrichment practices in commercial dairy farms. Some of the enrichment methods reviewed here may also be applied to those more extensive cattle raising systems, where similar challenges occur.

Key words: Social enrichment, Zero grazing, Animal welfare, Low resilience behaviours

## 1. INTRODUCTION

Nearly all dairy cattle are housed indoors, at least for some part of their life, and in an increasing number of farms, indoor housing is practiced all year round (van Vuuren and van den Pol-van Dasselaar, 2007; Winsten et al., 2010; March et al., 2014). In continuous indoor housing systems (also referred to as "zero grazing" systems), dairy cows are kept throughout the year in tie stall, free stall, or loose housing cowsheds. Access to pasture is either limited or absent. In the past, continuous indoor housing of dairy cows was practised mainly in regions where the climate was unsuitable for growing grass or too harsh for the animal. Nowadays, with the gradual shift towards intensified farming, year round housing is more widely practiced. It was recently estimated that zero grazing housing will become the most prevalent farming practice in North-West European countries such as NW-Germany and Denmark, by the next decade (Reijs et al., 2013). For example, in the Netherlands, the number of dairy cows housed indoors has tripled in the past 10 years (from 10% to 30%; CBS, 2015). In the United States, more than 95% of lactating cows are denied access to pasture (NAHMS, 2010). Other Mediterranean countries, such as Israel, now keep 100% of their dairy cows indoors throughout the year (Israeli dairy farms reform, 1999-2008). The practice of keeping cows indoors for extended periods may also result from environmental regulations aimed at reducing leaching of nitrates and phosphorus into water reserves (for example, the "Nitrates Action Programme" implemented in Northern Ireland from 2007; Nitrates Action Programme and Phosphorus Regulations 2015-2018).

Keeping animals indoors provides some important welfare benefits for the animals, such as protection from predators and toxic plants, reduced exposure to extreme weather conditions (Schütz et al., 2010) and external and internal parasites, as well as enabling the provision of a

64 nutritionally balanced diet throughout the year (Algers et al., 2009). However, it also confronts  
 65 animals with a wide range of challenges. These include abiotic environmental sources of stress  
 66 (e.g. exposure to loud and aversive sounds such as the noise produced by milking facilities,  
 67 Arnold et al., 2007, 2008; metal-on-metal clanging, Waynert et al., 1999) and confinement-  
 68 specific stressors that are more likely to be associated with indoor systems (e.g. restricted  
 69 movement when kept tied in their stall, when isolated at an early stage of life, or maintenance in  
 70 abnormal social groups, Morgan and Tromborg, 2007). Zero grazing systems, compared to other  
 71 production systems, are also associated with higher incidence of lameness (Haskell et al., 2006),  
 72 and increased risk for claw or foot problems, teat trampling, mastitis, metritis, dystocia, ketosis,  
 73 retained placenta and some bacterial infections (Algers et al., 2009). Once housed, animals are  
 74 forced to make substantial changes in their time budget (Newberry, 1993). For example, the food  
 75 searching and eating times of cattle may be reduced to 4 hrs a day, compared to 6-12 hrs on  
 76 pasture (Gomez and Cook, 2010) such that long periods of time must be occupied with a limited  
 77 range of possible behavioural patterns (Hughes and Duncan, 1988). Mason and Burn (2011)  
 78 argued that when the environment is too impoverished (i.e. without appropriate stimuli or  
 79 substrates) or too small, the ability of the animals to perform natural behaviours and to satisfy  
 80 their motivations (i.e. to fulfill their behavioural needs) is restricted. Such behavioural restriction  
 81 may result in frustration. Indicators of frustration in cattle include leg stamping (Cooper et al.,  
 82 2008; although this can also be associated with attempts to cope with forced standing by  
 83 alleviating strain on the legs and hoofs), non-nutritive oral behaviour (e.g. tongue rolling,  
 84 Ishiwata et al., 2007), and an increase in the visible percentage of eye whites (Sandem et al.,  
 85 2002; although the latter was also associated with fear, Sandem et al., 2004). Persistent  
 86 frustration is associated with the development of abnormal behaviours. One example is calves'

redirected oral behaviour toward pen mates, when fed from a bucket and restricted from performing suckling behaviour (Mason and Burn, 2011; Ninomiya, 2014).

Keeping animals in an environment that meets their proximate needs (“here and now”, Dawkins, 1983, such as feeding, drinking and sleeping) allows them to engage in low resilience behaviours (also referred to as 'luxury activities', i.e. behaviours that typically decrease when energy resources are limited or when the cost involved in the activity increases, McFarland, 1999), which are associated with improved welfare and long term fitness (Held and Spinka, 2011). One example is play behaviour, which drops out of the animal's behavioural repertoire in times of challenge (e.g. sickness, hunger, injury, predation risk and thermal stress). In the majority of cases, the presence of play behaviour is associated with improved welfare, and its disappearance is a reliable indicator of the transition from positive to poor welfare (Held and Spinka, 2011). In cattle, other low resilience behaviours include grooming (Borderas et al., 2008; Fogsgaard et al., 2012; but see also opposing findings by Almeida et al., 2008) and use of automated cow brushes (Mandel et al., 2013).

One strategy that can help animals to cope with stressors in their surroundings, prevent frustration and increase the fulfillment of behavioural needs is to enrich their environment. Newberry (1995) defined environmental enrichment as an improvement in the biological functioning of confined animals resulting from modifications to their environment. Biological functioning refers to increased fitness (i.e. lifetime reproductive success), increased inclusive fitness (i.e. indirect fitness, by helping genetically related individuals such as kin to increase their fitness), or a correlate of both such as improved health. By focusing on the biological functioning

of the animal, Newberry (1995) offered a practical and objective way to measure and evaluate the effect of different environmental enrichment methods on welfare. However, compromised welfare does not necessarily result *only* from impaired biological functioning (Fraser et al., 1997). For example, the welfare of bucket-fed calves is not reduced by malnutrition, but by an unfulfilled need to suckle (Fraser et al., 1997). For the purposes of this review, effective environmental enrichment will therefore be regarded as modification to the management or surroundings of the animal that demonstrably improves biological functioning (Newberry, 1995), or other validated measures of welfare (i.e. those measures that are correlated with valenced experiences, Nicol et al., 2009) over and above what is achieved by following minimum management standards (e.g. European Union guidelines).

Although environmental enrichment plays an important role in maintaining the wellbeing of zoo animals (Shyne, 2006), laboratory animals (Baumans and Van Loo, 2013), and certain livestock such as pigs (van de Weerd and Day, 2009; see also EU Directive 2008/120/EC), its implementation on cattle farms is limited and has not coincided with the gradual shift towards year-round indoor housing and the challenges it places on cows. Considering the global increase in the number of cows and calves raised in zero-grazing systems, exploring different methods for meeting their needs (e.g. by enriching their environment) is more relevant today than ever before.

This review has two aims; firstly, we will review recent evidence of the effect of environmental enrichment on the welfare of dairy cattle kept indoors. Secondly, this will be accompanied by an



assessment of the practicality of different enrichment options, considering the divergent needs of calves, heifers and cows separately when appropriate.

## 2. ENVIRONMENTAL ENRICHMENT

Bloomsmith et al. (1991) defined five categories of environmental enrichment; social, occupational, physical, sensory and nutritional. Each category may contribute to the welfare of the animal in a different way. For example, social enrichment may fill the need of calves for a companion and help them to cope better with stressors in their surroundings (de Paula Vieira et al., 2012), whereas physical enrichment in the form of providing a cow with a secluded area in an individual maternity pen may fulfill its need to hide when calving or when sick (Proudfoot et al., 2014a,b). When considering the contribution of each category to the welfare of the animal, it is important to note that: 1) The mechanism that underlies each category of enrichment, and each method of enrichment, can be multi-factorial (e.g. dam-rearing involves both social and nutritional factors, Kalber and Barth, 2014). 2) Each enrichment type can have both short- and long-term effects (e.g. drinking colostrum and milk ad-libitum is associated with improved weight gain at an early stage of life, but also aids in gastrointestinal-tract maturation, production of digestive enzymes and nutrient absorption at a later stage through hormonal factors found in the colostrum, Bach, 2012). 3) Each enrichment method can contribute differently to the welfare of the animal, when applied at different stages of the production cycle, as a calf or an adult cow (e.g the presence of a conspecific is associated with improved cognitive development in calves, Gaillard et al., 2014; but also with reduced stress during social isolation in cows, Kikusui et al., 2006). 4) The contribution of each category to the welfare of the animal can be explained by more than one mechanism (reduced stress expressed by calves that were raised in pairs,

compared to singly reared calves, can be explained both by improved cognitive development, which allows better learning abilities but also by the mere presence of a conspecific (social buffering)). 5) Some enrichment methods are limited to a certain time window (e.g. the effect of early social housing on cognitive development, Gillard et al., 2014). The examples mentioned above, and others, will be discussed in more details throughout the text and will serve as the first step towards developing a conceptual framework for enriching the environment of dairy cows. Such a framework could allow better understanding of cattle needs and improve the implementation of enrichment practices in dairy farms.

## **2.1 Social Enrichment**

Social enrichment is defined as the provision of access to direct or indirect (visual, olfactory, auditory) contact with conspecifics (other individuals of the same species) or humans (Bloomsmit et al., 1991).

On most North American and European dairy farms, calves are separated from their mothers within 24 hours of birth (Algers et al., 2009; but see also dam and foster rearing in organic farming, Kalber and Barth, 2014). Separating the dyad is stressful for both the dam and her calf and followed by behavioural (e.g. increased calling and activity; Weary and Chua, 2000; Flower and Weary, 2003) and emotional (i.e calves: cognitive bias test; Daros et al., 2014) responses. When carried out within 24 hours of birth, the process is accompanied by a weaker behavioural response compared to separation at a later stage (i.e. two weeks after birth; Flower and Weary, 2001; but see Hudson and Mullord, 1977 for behavioural distress following 5 minutes of calf-dam contact, and Johnsen et al., 2015 for the formation of cow-calf bond without nursing). Early

separation is also done for economic (e.g. use of sources of nutrition that are cheaper than feeding calves' with their mother's milk, such as milk replacers or 'waste' milk that cannot be sold) and management reasons (e.g. close monitoring of calves health and food consumption; minimizing the transmission of Johne's disease, Collins et al., 2010, and faster milk-let-down speed, Kilgour and Dalton 1984). Following separation from the dam, calves are kept in single housing, pair housing or in group housing (three calves or more).

### ***2.1.1 Contact with Conspecifics***

**Calves.** In farms that practice single housing, calves are reared individually in pens or hutches for two to eight weeks, mainly with the aim of decreasing the risk of horizontal disease transmission, but also for helping farmers to monitor calf milk intake and health. The association between group housing and morbidity is affected by the size of the group, with calves kept in large groups (i.e. 7 or more) being at higher risk of disease (Losinger and Heinrichs, 1997, Wells et al., 1997; Svensson et al. 2003, Svensson et al., 2006; Svensson and Liberg, 2006). A tradeoff between single housing and group housing is pair housing, where physical contact is limited to only one other calf, and the risk of pathogen transmission is limited. A recent study compared the effect of different levels of social contact on calf health ranging from strict isolation in single housing to full physical contact in pair housing, but found no effect of degree of social contact on the level of the 5 most common pathogens present in Danish calf feces, or on the development of serum antibodies against the three most common respiratory pathogens (Jensen and Larsen, 2014). However, in indoor environments with poor ventilation and drainage, keeping calves in small groups (i.e. three calves per pen) or in pairs may increase the risk of respiratory disease, compared to individual rearing (Cobb et al., 2014). Acknowledging the contribution of physical

contact to disease transmission, pair housing can be considered a tradeoff between individual rearing and group housing, which allows calves to engage in social contact, while limiting disease transmission.

The need of calves for social contact with their peers is present from the first week of life (Wood-Gush et al., 1984). When given the option, they are more motivated to get access to full physical contact with their conspecifics, compared to only head contact through metal bars (Holm et al., 2002). Calves raised with full social contact (either from birth or 3 weeks of age) will establish stronger bonds with their group members, compared to calves raised with limited contact (Duve and Jensen, 2011). The bonds that calves develop at an early stage will affect their social preferences as adults (Sato et al., 1993; Færevik et al., 2006; Raussi et al., 2010; Gygax et al., 2010).

Physical contact with conspecifics from an early age affects calf development; individually housed calves, compared to paired housed ones, spend less time at the feeder, visit it less frequently and start ingesting concentrate from computerized starter feeder at a later stage (de Paula Vieira et al., 2010, see also Warnick et al., 1977 and Hepola et al., 2006). Individual rearing also reduces calves' social skills and their ability to cope with environmental stressors (de Paula Vieira et al., 2012; Jensen and Weary, 2013). They are also more fearful of unfamiliar calves (Jensen and Larsen, 2014; de Paula Vieira et al., 2012), have a higher heart rate during confrontation (Jensen et al., 1997), struggle more when restrained for blood sampling (Duve et al., 2012) and vocalize more when weaned from milk (de Paula Vieira et al., 2010). The reduced vocal response of pair-housed calves when weaned is suggested to reflect the effect of social

buffering (i.e. the alleviation of stress responses attributed to the presence of a conspecific, Edgar et al., 2015), which can help modulate or down regulate the impact of stressors on the homeostasis of the recipient (for a recent review, see Rault, 2012). Indeed, calves that were isolated with no companion for a period of 20 minutes, were found to vocalize more, show less locomotor activity and explore the pen less, compared to those that were isolated with a companion, particularly a familiar companion (Færevik et al., 2006). Social isolation at an early stage seems to also affect calf behaviour later on in life. Broom and Leaver (1978) studied the effect of prolonged social isolation (8 months) on calves' behaviour at the age of 8 months and 20 months. In both age groups, individually housed calves spent more time alone and had a lower social rank once introduced into a new group, compared to grouped housed calves.

An additional mechanism to social buffering that can affect the behaviour of calves raised in isolation is impaired cognitive development. Individually reared calves achieved poorer performance in a colour discrimination reversal learning task than calves reared in pairs (Gaillard et al., 2014). The socially reared calves appeared to be more flexible in their response to change in routine management and housing, an ability that was previously associated with improved welfare (Wechsler and Lea., 2007). One implication is that socially-reared animals may be more competent in interacting with new technologies, such as robotic milking equipment and automated feeders. Indeed, de Paula Vieira et al. (2010) showed that calves raised individually were slower at learning to use a computerized starter feeder compared to calves raised in pairs. Therefore, the reduced learning ability associated with individual rearing may result from a cognitive impairment or emotional deficiency (or both). Raising calves in pairs or small groups

fulfills their need for social contact with conspecifics from an early age, help to develop cognitive skills, social skills, and reduce stress-associated reactions.

Pair housing also has economic benefits. Rearing calves in pairs requires less space than individual housing, which can be used for spacing the pens further apart (i.e. lowering the chance of horizontal disease transmission), and for increasing the living area for each pair to allow greater comfort and encourage play behaviour. For example, in order to raise 10 calves in isolation (0.9\*1.8m per calf, with 1 meter space between pens), farmers need an area which is 18 meters wide. In the same area, using pair-housing pens, farmers can increase the living space of each calf by more than 35% (1.35\*1.8m per calf instead of 0.9\*1.8m) and the distance between pens by more than 10% to minimize further the risk of diseases transmission. Housing calves in pairs however, requires adjustment to the feeding method to minimize food competition and decrease cross-sucking behaviour (addressed below under nutritional enrichment).

Although pair housing is a promising rearing solution that balances calf health and social needs, some fundamental questions regarding the timing of its implementation still remain open. Other questions concern the implications of dyad separation at a later stage of life (i.e. for either short periods of time, e.g. for a husbandry procedure or for long periods of time, such as when kept in different feeding groups). Breaking a social bond between two calves raised together from the first day of life (and prevented from maternal contact) may prove to be as stressful as breaking the bond between a calf and its dam bonded for a similar amount of time. To our knowledge, these questions have not been addressed yet and demand further investigation. Better understanding of calf social needs (i.e. especially in early life, when kept in isolation) will allow

us to integrate more carefully their basic health and functioning, affective states and natural living (Fraser, 2008).

A more natural rearing method that is little practiced in intensified dairy farms, is to keep calves with their dams following parturition. Dam reared calves are either kept with restricted or full contact with their dam/foster cow and often have access to other calves and adult cows. The length of the rearing period may vary from days to months according to farm management. Calves reared with their dam, compared to calves housed in groups and fed from an automatic feeder, express less abnormal oral behaviour (e.g. cross suckling). Interestingly, low cross suckling rate was documented in both restricted (i.e. twice a day for 15 minutes each) and unrestricted contact with the dam (Roth et al., 2009; Hillmann et al., 2012). Dam-reared calves also struggle less when restrained for blood sampling compared to those housed singly or in pair (Duve et al., 2012). When submitted to an isolation test, calves that have been reared with their dam show lower increase in salivary cortisol concentrations compared to artificially reared calves after reunion with their dam/group (Wagner et al., 2013). Rearing calves with their dam/foster cow also seems to affect their behaviour later on in life (Le Neindre, 1989; Wagner et al., 2012; 2015). When faced with the challenge of integration into a new group, dam reared heifers (either with restricted or unrestricted contact) express more submissive behaviour associated with longer duration of feeding and earlier lying activity, compared to heifers that were separated from their dam and fed through an automatic feeder (Wagner et al., 2012). In another study, 2.5 year old cows that had permanent access to their dams during the first 12 weeks following parturition, expressed lower sympathetic and higher HPA-axis reactivity compared to cows that were fed by an automatic feeder (Wagner et al., 2015). The latter finding

suggests that calves reared with their dam develop a reactive coping style later on in life (Wagner et al., 2015). However, since the efficacy of the coping style (i.e. reactive and proactive) depends on the situation/environment, the welfare implications of this finding are not yet clear.

Keeping calves with their dam is regarded to be a natural rearing method that benefits from better public opinion compared to methods that involve early separation (Ventura et al., 2013). However, concerns regarding cow health (e.g. transmission of Johne's disease through contact of calves with the feces of its dam), and impaired milk ejection still exist (Kalber and Barth, 2014). Indeed, the opponents for this rearing method base their arguments on the possible negative effect on calf and cow health, as well as the emotional distress that will be caused once the cow-calf bond is broken later on, and the limited ability of the industry to accommodate cow-calf pairs (Ventura et al., 2013). In dairy farms, which are free of Johne's disease and can manage the logistics that are associated with keeping the dyad together (e.g. suitable enclosures, clean and dry environment), rearing calves with their dam could be an alternative enrichment method to pair housing that benefits from better public opinion. An alternative option that favors one side of the calf-dam dyad is to raise calves with a foster cow. The latter is suggested to allow calves to satisfy their suckling motivation and engage in social contact with adult cows, and may reduce weaning stress (Kalber and Barth, 2014). However, more knowledge about this system is needed in order to evaluate its contribution to the welfare of calves and its possible negative effect on the welfare of the foster cow (e.g. once more than two calves are fed from the same cow).

**Cows.** Cows are grouped based on their physiological status (lactating/dry) or milk production status (low vs. high milk yield). As their status changes, they are regrouped and must form



relationships with the new group members. This can be stressful. For example, regrouping destabilizes the social dynamic within the group and increases physical competition in the hours and days following regrouping (von Keyserlingk et al., 2008). Indeed, cows that enter a new group experience increased displacements from the feeding area and their eating time, lying time, number of lying bouts and allogrooming events are reduced. In addition, milk production is reduced on the first day after regrouping (von Keyserlingk et al., 2008). Basic husbandry practices, such as reducing the stocking density in the pen (Talebi et al., 2014), using a familiar pen for regrouping the cows (Schirmann et al., 2011), or other methods such as grouping during the evening hours (compared to mornings, Boyle et al., 2012), can help lessen the negative effects of this procedure.

The ideal solution to meet the social needs of the cows would be to keep them in stable groups. This would therefore allow them to enjoy the benefits of social companionship, and to benefit from social buffering, enabling better coping with stressors (Gutmann et al., 2015). The efficacy of social buffering depends on the degree of affiliation between the interacting partners (calves: Færevik et al., 2006; cows: Gutmann et al., 2015, for bulls see Mounier et al., 2006). Social buffering in cattle can be achieved via grooming behaviour (i.e. licking), which depends mainly on familiarity and increases with the length of cohabitation (Sato et al., 1991). This behaviour is regarded as a reliable indicator of friendship (Boissy et al., 2007), and seems to be independent of social dominance, as solicitation occurs both from dominant and subordinate cows (Sato et al., 1991; Val-Laillet et al., 2009). In addition to helping cows to stay clean (i.e. remove parasites, Guillot, 1981), it is suggested that this behaviour induces a "physiological calming effect" (Laister et al., 2011) and helps to resolve conflicts (Val-Laillet et al., 2009). Licking behaviour in

cows reduces the heart rate of the receiver (Laister et al., 2011), and was found to be directed more towards lame cows compared to non-lame cows kept at the same stall (Galindo and Broom, 2002). These findings suggest that licking behaviour may have a role in alleviating discomfort (Galindo and Broom, 2002). In cases where grouping is necessary, an intermediate solution could be to regroup cows in the company of familiar conspecifics in order to promote such affiliative interactions (Bøe and Færevik, 2003). Heifers that are introduced to a herd with a familiar conspecific (i.e. in pairs) face significantly less agonistic interactions compared to singly introduced heifers ( $7.19 \text{ h}^{-1}$  vs.  $3.79 \text{ h}^{-1}$ , Neisen et al., 2009) and integrate faster into the herd (Gygax et al., 2009). Heifers that are introduced to a new group in pairs show greater resemblance between their time budget and the time budget of other cows in the herd (e.g. time spent in the lying areas and feeding areas), compared to heifers that were introduced singly (Gygax et al., 2009, for increased lying times of heifers integrated in pairs compared to singly introduced heifers, see also O'Connell et al., 2008).

### ***2.1.2 Contact with Humans***

When kept indoors, cows rely on humans for almost every aspect of their lives. The interaction with humans is at times, inevitable, and varies between farms according to management policy, the size of the herd and the level of automation in the farm. Daily interaction with humans has a significant effect on cows' behaviour and productivity (Hemsworth, 2003). Humans can evoke fear in animals by virtue of their relative size, and their propensity for quick or unpredictable movements (Rushen et al., 1999). For example, the use of negative interactions by stock people (i.e. slaps, pushes, or hits with the hand or an object such as a plastic pipe) are negatively correlated with milk yield, protein, and fat, as well as with increased flight distance of cows (i.e.

the percentage of cows approaching within 1 m of an experimenter in a standard test; Hemsworth et al., 2000). However, in order for a cow to experience a negative interaction with a human, engaging intentionally in negative handling is not mandatory. Several of the routine husbandry procedures practiced in dairy farms are intrusive by nature and can be aversive to cows, causing pain and/or stress (Pilz et al., 2012). For example, artificial impregnation of cows involves a series of intrusive procedures, which starts with the artificial insemination itself (i.e. including both rectal and vaginal intrusions) and continues with routine pregnancy examinations that involve rectal examination. Following an aversive treatment, dairy cows and calves learn to avoid the specific handler and place associated with the aversive experience (“learned aversion”, Taylor and Davis, 1998, de Passillé et al. 1996; Rushen et al., 1998). By practicing positive handling from an early age, farmers can both help their animals to reduce stress responses during aversive experiences (in the case of artificial insemination, positive interactions by stockpersons was found to be positively correlated with conception rates, Hemsworth et al., 2000) and induce positive affective states promoting positive welfare in the farm (Ellingsen et al., 2014; Proctor and Carder, 2014; 2015a,b).

**Calves.** In calves, positive handling at 4 weeks of age (e.g. moving slowly and calmly around in the pen, speaking in a quiet and calm voice and encouraging interactions including pats and scratches) affects approach behaviour towards familiar handlers, compared to negative handling (e.g. fast movements, speaking with a harsh voice and creating noise with different tools, such as a plastic bottle with rattling stones in it ;Schütz et al., 2012). However, the impact of handling quality at an early age may be overshadowed by the amount of human contact at a later stage (after 3 months, Schütz et al., 2012). When test calves (handled negatively and positively) were

compared with a control group reared routinely on-farm (i.e. without any particular handling treatment), the latter showed greater avoidance behaviour than both positively and negatively handled calves, demonstrating the importance of both quality and quantity of handling. In a more recent study, gentle interactions on group-housed dairy calves during the first 14 days of life were associated with smaller avoidance distances from a familiar person before and after weaning (i.e.  $86.2 \pm 5.1$  days old), and with higher average daily gain (up to 7%, depending on milk provision, Lürzel et al., 2015). In beef cattle, gentle interactions (i.e. 120 min of touching over 6 days within 4 weeks of life) were associated with less fearful behaviour towards humans (i.e. smaller avoidance distances) and less stress-related behaviour at the abattoir at the age of 10 months (i.e. less avoidance behaviour in the stunning box and a tendency for lower cortisol concentrations in the blood taken during exsanguination, Probst et al., 2012). Positive handling is also associated with more positive mood, as assessed by qualitative behaviour assessment (QBA, Ellingsen et al., 2014). In their study, Ellingsen et al. (2014) characterized the behaviour of stockpersons and portrayed the body language of the dairy calves that were under their care. The stockpersons who handled their calves patiently, and petted and calmly talked to them during handling, were rated by respondents to be more 'friendly' and 'content'. By contrast, stockpersons with a nervous handling style, or who were dominating and aggressive had calves that were perceived as in a more negative mood (Ellingsen et al., 2014). However, the extent to which these QBA labels reflect real underlying mood states needs to be validated using more direct measurements of animal behaviour and physiological state (e.g. Mendl et al. 2010). The effect of brushing calves by humans will be reviewed under sensory enrichment.

**Cows.** In cows, a high percentage of positive interactions (talking quietly, petting and touching) and a low percentage of negative interactions (forceful use of stick or hand, shouting and impatient talk) are associated with reduced avoidance of handlers in the milking parlor (Waiblinger et al., 2002). Waiblinger et al. (2004) found that the stress response to an aversive veterinary procedure could be reduced by positive handling prior to, and gentle interactions during the procedure. In this study, positive handling included feeding cows a small amount of concentrate out of the hand, stroking them at the neck and head as long as they accepted it, and speaking to them in a soothing way for several minutes during 10 days distributed over a 4 week period. Gentle interactions applied during an aversive veterinary procedure consisted of stroking the cows at the neck and head and speaking to them in a soothing way. The authors found that cows previously handled in a positive way had a lower heart rate and kicked less when isolated for the procedure. In addition, cows that were calmed down during the procedure itself (i.e. gentle interaction) expressed less restless behaviour (e.g. head shaking and tail flicking). The success of "calming down" the animals however, differed between handlers (Waiblinger et al., 2004; for additional evidence of the effects of positive and negative handling, see Rushen et al., 2001; Lensink et al., 2001; Waiblinger et al., 2006; Breuer et al. 2000 and Hemsworth et al. 2000; but see also Pajor et al., 2003 and Stewart et al. 2013 for limited or no effect of previous handling). Brushing by humans can also lower stress levels in times of isolation and will be discussed later under tactile enrichment.

## ***2.2 Occupational Enrichment***

Occupational enrichment encompasses both enrichment that encourages exercise and psychological (also referred to as cognitive) enrichment (e.g. devices that provide animals with control or opportunities to use their cognitive abilities, Bloomsmith et al., 1991).

### ***2.2.1. Enrichment that Encourages Exercise***

**Calves.** During early stages of life, calves are raised alone, in pairs or in groups. The space allowance for each calf varies between rearing systems (and stocking density), and has a direct effect on their ability to engage in voluntary physical training (Jensen et al., 1998). Physical training of animals can be achieved through play behaviour, which can be directed at a conspecific, object, conducted alone or in a company (Held and Spinka, 2011). In calves, play behaviour was studied mainly in rearing conditions that allow very limited voluntary movement (i.e. confinement). The motivation to perform locomotor play (i.e. galloping, bucking and kicking, which involves no interaction between individuals) and trotting, was shown to increase with the length of confinement (Jensen, 2001) suggesting that play is a behavioural need. A reduction in play behaviour in young mammals has been proposed as a reliable indicator of the transition from positive to poor welfare (Held and Spinka, 2011). Indeed, play behaviour in calves is shown to decrease when food provision drops (Duve et al, 2012), or when calves are subjected to painful procedures (e.g. hot-iron disbudding, Mintline et al., 2013). Improving the welfare of calves by rearing them in groups is associated with higher rate of play behaviour (Jensen et al., 1998). In a recent study, Valníčková et al. (2015) assessed the effect of social companionship (single vs. group housing) on play behaviour. In this study, housing calves in groups was associated with more spontaneous play behaviour in the pen compared to housing

calves individually. In contrast, in an open-field test and during a social test (i.e. encounter with an unfamiliar conspecific), individually housed calves were more playful compared to group housed calves (Valníčková et al., 2015). The lower levels of spontaneous play and the higher rebound effect during the open field test and during the social test (i.e. when calves were exposed to a larger space without or with companion), was suggested as indicating deprived natural levels of play behaviour in individually housed calves (Valníčková et al., 2015). Although the occurrence of play behaviour is generally low, its applicability as a measure of welfare in combination with other indicators is worth investigating further. Additional methods for encouraging exercise (i.e. via play behaviour) will be further addressed under physical enrichment.

**Cows.** The motivation of cows to engage in physical activity (i.e walk, trot, gallop and jump), has mainly been studied in individuals kept in tie stall barns. Despite increasing criticism, tie stalls are still used throughout the world, with over 70% of dairy farms in Canada (Canadian Dairy Information Centre., 2014), 38% in the United States, 69% in Finland, 78% in Switzerland (Barkema et al., 2015) that tether their cows. In the EU, 20% (lowland) and 80% (upland) of cows are tethered at least during the winter (Veissier et al., 2008). Cows, similarly to calves, show increasing motivation to engage in physical activity as a function of the time they spend in confinement (i.e. tied up, Loberg et al., 2004; Veissier et al., 2008). Daily access to an exercise area (e.g. indoor exercise area, out-door paddock, or preferably pasture) has been shown to revert locomotor activity of tied cows to levels observed in loose-housed cows (Veissier et al. 2008), and provides them with more opportunities to engage in social interactions, explore their environment and groom 'hard to reach' hindquarters (Krohn et al., 1994). In Hérens cows (i.e. a

Swiss breed that is highly motivated to engage in dominance interactions), daily access to an exercise yard was associated with lower frequency of agonistic interactions compared to access every 3 days or more (Castro et al., 2011). Likewise, Loberg et al. (2004) found lower levels of aggressive behaviours (i.e. threatening and pushing) in tied cows that had access to an exercise yard once a day, compared to once a week. Engagement in exercise is also associated with positive effects on claw conformation (Loberg et al., 2004), and reduced incidence of lameness (Regula et al., 2004) and of mastitis (Popescu et al., 2013). In farms where tethering is still practiced, daily exercise of more than one hour is recommended as a measure to fulfill, at least to some extent, the motivation of cows to engage in movement (Veissier et al. 2008). For minimizing hock lesions, at least 50 hr per month are recommended (i.e. preferably on pasture to allow the cows to lie down comfortably, Keil et al., 2006). In the latter study, the duration of each exercise period seemed to be particularly important, as increased frequency alone was not found to be beneficial. To our knowledge, the contribution of additional daily exercise to the welfare of cows kept solely in free-stall systems (i.e. which can be crowded, with high traffic areas, and walking surfaces that are very difficult to keep dry and are slippery) has not yet been studied, and its practicality for high yielding dairy cows and for calves should be considered separately.

### ***2.2.2 Cognitive Enrichment***

Recent studies have shown that farm animals are capable of more complex cognitive and emotional responses than previously thought (Broom, 2010). Yet, farm animal housing generally offers less stimulation and fewer opportunities for animals to use their cognitive abilities than those available in the wild (Langbein et al., 2009). Providing animals with more opportunity to



use their cognitive abilities has been suggested to be an important component of animal wellbeing (Carlstead and Shepherdson, 2000), however, this key assumption still requires additional supporting evidence. Cognitive enrichment can give animals control over aspects of their environment, and can lead to positive affective states. The majority of empirical work on the effects of providing animals with control over their environment has investigated control over punishment rather than reward (Sambrook and Buchanan-Smith, 1997). However, in recent years, there has been increasing evidence suggesting that the control of positive situations, such as situations that involve anticipation of consummatory reward, improves welfare (Manteuffel et al., 2009; Basset and Buchanan-Smith, 2007).

Manteuffel et al. (2009) suggested that cognitive enrichment of group-housed animals on commercial farms could be achieved using self-controlled operant learning tasks, and adapting the degree of challenge to the cognitive abilities of each species. The initial stress and frustration which may arise when a challenge is presented to the animal is suggested to be an important feature in the process of cognitive enrichment, as long as the animals possess the skills and resources to effectively solve the problems that they face (Meehan and Mench, 2007). Habituation and "over-experience" however, should be prevented by changing, to some extent, the conditioned discriminatory stimuli or by adding a further conditioned behaviour (e.g. variable or fixed ratio lever pressing) to the initial one. In recent years, an increasing number of studies have explored, both directly and indirectly, the cognitive abilities of cattle (e.g. face perception and recognition, Coulon et al., 2011; spatial memory and decision making, Bailey et al., 1989a, Bailey et al., 1989b; operant conditioning and reversal learning, Vaughan et al., 2014; Webb et al., 2014; Wechsler and Lea 2007; Wredle et al., 2006; reversal learning in calves, Gaillard et al.,

2014). Nevertheless, to our knowledge, only Hagen and Broom (2004) specifically explored the effect of cognitive enrichment on cattle behaviour. In this study, heifers were divided into an experimental group, where heifers were conditioned to press an operant panel to open a gate and gain access to a food reward, and a control group, where the gate opened automatically after a delay equal to their matched experimental group partner's latency to open it. Heifers that learned the operant task displayed evidence of greater "excitement" (i.e. higher heart rates and more vigorous movement), compared to control heifers, which received the same reward after spending the same amount of time in the pen. However, it is difficult to infer from this study the valence associated with the higher arousal of the experimental group, and thus whether these heifers indeed experienced enjoyment (contingent on understanding and gaining control over the task) or frustration (Spinka and Wemelsfelder, 2011).

When planning cognitive enrichment for a group of cows, individual differences arising from age or previous experience must be considered (Manteuffel et al., 2009). Indeed, individual differences can make an enrichment task solvable and rewarding for one group member, and difficult and frustrating for another. Setting the complexity level of a task to the ability of the weakest animal in the group will allow all of the animals in the group to solve the task, but may be too simple and easy for the majority of the group. A more promising solution that has, to our knowledge, not been tested yet in farm animals, would be to adjust the complexity level of the task to each member of the group. A system based on machine learning algorithms could adjust itself to the (changing) lifetime abilities of each individual. This kind of knowledge could be added to the individualized data that is already being collected in an increasing number of dairy

farms around the world (e.g. daily milk yield, daily activity or rumination time), and give the farmers a more detailed image of each individual animal kept in the group.

### **2.3 Physical Enrichment**

The complex cognitive abilities of cows should also be considered when designing housing and husbandry systems (Broom, 2010). Physical enrichment includes altering the size or complexity of the animal's enclosure or adding accessories to the enclosure such as objects, substrate, or permanent structures (Bloomsmit et al., 1991).

**Calves.** Providing animals with access to alternate enclosures (i.e. by dividing space into different functional areas) was suggested to increase opportunities for exploration and patrolling, as well as opportunities for camouflage and hiding (Newberry, 1995). Dividing the space within an enclosure may also be beneficial for decreasing antagonistic interactions between calves. Ninomiya and Sato (2009) compared the rate of agonistic encounters (e.g. head butting, chasing, escaping) between Japanese black calves kept in an enriched pen divided by a wooden wall, with the rate of agonistic encounters between calves kept in a control, not divided pen. Dividing the space in the pen resulted in lower motivation of stronger calves to chase the weaker ones, possibly by preventing eye contact between the animals and forcing them to pass through narrow gaps in order to come into contact with each other (Ninomiya and Sato, 2009). However, since the authors did not control for additional factors, such as the effect of the brush and wood log that were also placed in the enriched cage, the role of the wooden wall in decreasing agonistic behaviour cannot be clearly demonstrated.

Alternating the physical environment of calves, in addition to reducing agonistic behaviours, can be used to encourage play and related exercise. Providing calves with increased space allowance (i.e. 1.8m× 3.0m compared to 0.9 m × 1.5m) is associated with higher levels of play behaviour (Jensen et al., 2015a). Adding additional stimuli to the enclosure (i.e. provision of fresh bedding) also appears to stimulate the occurrence of this behaviour (Jensen et al., 1998, see also Schütz et al., 2012). Jensen et al. (1998) reported that provision of straw in connection with morning feeding (i.e. when fresh straw was supplied every day after milk feeding), stimulated the largest daily peak in play behaviour. Playing with straw was reported by the authors to resemble ground play, as opposed to object play. Ground play consists in the calf rubbing the head and neck against the ground while kneeling down (Schloeth, 1961). However, since fresh straw was always supplied following morning feeding in Jensen et al. (1998), its effect on play behaviour cannot be fully evaluated, as it is not possible to disentangle time of day effects from the effect of straw. Encouraging play behaviour in calves by the use of other substrates or external objects (e.g. cow brushes and/or balls hanged at a height of 1.3 meters) may prove to be beneficial when social isolation is mandatory (e.g. quarantine, Bulens et al., 2014).

**Cows.** In zero grazing systems, cowsheds are usually designed to provide constant visual and physical contact between conspecifics. Allowing cows to maintain contact with each other, corresponds, at least to some extent, to their natural need to live in a group. However, in the wild, cows also have the possibility to isolate themselves from the group when they need to, such as the time around calving (Lidfors et al., 1994). The need to isolate oneself from the group seems to be important also for the high yielding cows used today in the industry. Proudfoot et al. (2014a) showed that, when given the opportunity, dairy cows housed in individual maternity

pens preferentially used a secluded area to calve. Cows began using the secluded area more than usual during the hour before calving and continued to use it more for the hour after calving. The need to isolate oneself from the group was also documented during times of illness. Proudfoot et al. (2014b) found that cows with high rectal temperature after calving and signs of an infectious disease (mastitis, metritis, pneumonia, or some combinations of these diseases) spent more time in the secluded area compared to healthy cows. Building secluded areas inside the cowshed thus seems to provide cows with the opportunity to express, at least to some extent, their natural need to isolate themselves from the group. Cows should have free access to these secluded areas, as the need for isolation at times of morbidity is not shared by all group members, and not for all morbidity cases (e.g. lame cows, Jensen et al. 2015b). Secluded areas that cows can access freely can be introduced in both loose housing systems (e.g. by installing a "movable" fence that can be folded when the soil needs to be cultivated), and in free stall systems (e.g. by installing dense nets that provide a visual barrier but can still allow air flow). Monitoring the occupation of such areas (i.e. in calving pens, hospital pens and free-access secluded pens) may prove to also be beneficial for farmers, as they could acquire an additional indicator for the wellbeing of their cows and their proximity (i.e. in time) to calving. The need to minimize contact with the group can also appear in low ranking dairy cows at times of high aggression in the shed (feeding time, Haskell et al., 2013). For example, providing access to a loafing area allows low-ranking cows to avoid dominant animals (Haskell et al., 2013). The extent to which summer yards/exercise yards (i.e. adjacent to the cowshed that allow animals to "sun bath") can further contribute to enrich the physical environment of cows should be explored in future studies.

Enriching the lying area of cows by providing them with a comfortable lying surface (i.e. soft, non-abrasive, clean and dry) was also suggested to be an important factor in enhancing cows' comfort and welfare (Tuytens, 2005, Fregonesi et al., 2007). Bedding materials can vary from organic or mostly organic materials such as straw, saw dust, inorganic (e.g. sand) to synthetic materials such as or mats made of rubber combined with polypropylene and nylon. The preferences of cows can strongly differ depending on the type of bedding (Norrington et al, 2008), quantity of bedding (Jensen et al., 1988), quality of bedding (i.e. whether the lying surface is dry or wet; Fregonesi et al., 2007), on the season (summer and winter, Manninen et al., 2002), and on previous experience (Tucker et al., 2003). However, their preference for some bedding materials may not necessarily fit the long term benefit of the animal. For example, bedding with straw, compared to sand bedding, was associated with increased lying time (straw  $749 \pm 16$  vs. sand  $678 \pm 19$  min per day). However, it was also associated with more severe hock lesions and with less improvement in overall foot health (Norrington et al, 2008). The costs of bedding materials can be high in some countries and therefore, lying mats can also be used as a measure to reduce the costs of bedding, by placing small quantities of bedding materials on them (Norrington et al., 2010; for further information on bedding, see review by Tuytens 2005). Other measures that can enrich the physical world of cows, such as placing cow brushes inside the cowshed, will be addressed in the sensory enrichment section.

#### ***2.4 Sensory Enrichment***

Sensory enrichment is defined as stimulation designed to trigger one or more of an animal's senses; Wells, 2009). The stimulation can be achieved through visual (e.g., television), auditory (e.g. music, vocalizations), or other modalities (e.g., olfactory, tactile, taste; Bloomsmit et al.,

1991). As there are very few studies conducted in each subfield of sensory enrichment, we shall discuss both age groups (i.e calves and cows) together.

#### ***2.4.1 Auditory Enrichment***

When considering the use of auditory enrichment, the excellent hearing of cows, compared to humans, should be taken into account. The hearing range of cattle lies between 23 Hz and 35 kHz, nearly one octave higher than that of humans (Heffner and Heffner, 1983). Exposing cattle to high pitch sounds was suggested to damage their hearing and affect feeding behaviour (Johns et al., 2015). The environment on commercial dairy farms can be loud and noisy; the sound of metal gates opening and closing, tractors carrying total mixed ration/straw and vacuum operated milking machinery, are only some of the noises that cows are exposed to frequently. Waynert et al (1999) reported that beef heifers exposed to the noise of clanging metal and humans shouting showed higher heart rate and activity than cattle in a quieter environment. Similarly, noise recorded from a commercial milking facility induced fear in dairy heifers and, when given the opportunity, the heifers learned to avoid these noises (Arnold et al., 2008). Although the results of the latter study cannot be generalized to all kinds of milking machinery, and may depend on farm management, they align with the argument that an overall reduction of ambient noise, rather than additional acoustic stimulation, might be more important for the animals themselves (Wells, 2009). In accordance with this view, Newberry (1995) emphasized that adding auditory stimuli to an environment that is already noisy may cause more harm than good, especially if the animals have no control over the sound (i.e. such as the ability to move to a quieter location or to switch off the sound).

On farms, auditory stimuli are used for a variety of tasks, from improving cow traffic in automatic milking systems (Uetake et al., 1997), to training cattle to approach a food source (Wredle et al., 2004). In most cases, the association between the auditory stimulus and cow behaviour is established using classic or operant conditioning. In a recent study, Kızılcı et al. (2013) used music as a measure to reduce cows' stress levels during milking. Cows that were played classical music (during the milking period, for a period of 28 weeks), compared to no music, had higher milk let down speed ( $6.27 \pm 0.12$  min vs  $6.68 \pm 0.13$  min, respectively). However, since the authors did not control for the effect of classical music on the workers and its possible effect on the handling of cows, nor did they measure stress levels in the cows, the association between classical music and stress levels cannot be clearly determined. If indeed playing classical music in the milking parlor has an indirect effect on the welfare of cows (i.e. by affecting the milkers themselves), an intermediate solution that would reduce ambient noise for cows while enabling farmers to hear music would be the use of earphones.

In addition to classical music, some studies tested for the effect of more “natural” auditory stimuli, such as playbacks of calf vocalizations, on increased milk yield. Pollock and Hurnik (1978) exposed cows in the milking parlor to playbacks of calf calls. The broadcast of calls started just as the cows entered the milking parlor and ceased when the last teat cup had been placed on the last cow. Milk let down during the first two minutes of milking was higher in the treatment group compared to the control group, which was not exposed to calls (average production of 16 cows =  $81.1 \pm 0.4$  kg vs.  $79.7 \pm 0.5$  kg, respectively). In a more recent study, McCowan et al. (2002) reported an increase of 1-2% in milk yield during the second milking session following playbacks of calf calls. Although the results of these two studies present a



positive association between exposure to playbacks of calf vocalizations and milk yield/release, their effect on the emotional state of the cow was not assessed (McCowan et al., 2002). It has been suggested that, similarly as in most other mammals (Manteuffel et al., 2004; Briefer, 2012), vocalizations in cattle may signal the physiological and emotional state of the calling animal (Thomas et al., 2001; Marchant-Forde et al., 2002; Ikeda and Ishii, 2008). Indeed, previous studies have shown that calves vocalize more before feeding time and that calves fed by conventional management (i.e. twice daily for a total of 5 L during 24 h) produce more calls compared to calves fed every 4 hours with 8 L of milk per day ( $31.4 \pm 7.0$  vs.  $5.0 \pm 3.4$  respectively; Marchant-Forde et al., 2002). The calls used in the two above-mentioned experiments (Pollock and Hurnik, 1978 and McCowan et al, 2002) were produced by calves that were either deprived of food (for 8.5 h, Pollock and Hurnik, 1978) or prior to milk feeding (McCowan et al, 2002). Marchant-Forde et al. (2002) found that cows that were played recordings of calves (i.e. produced in similar conditions - in a commercial farm, under one week of age, before feeding), compared to white noise, had greater heart rate response, increased head movements and ear movements, oriented more toward the speaker and spent less time eating. Since calf call playbacks were played to the cows throughout the milking session in Pollock and Hurnik, (1978) and McCowan et al, (2002) (i.e. playback calls did not stop while milking was in process), a negative emotional reaction in the cows might have been triggered through, possibly, emotional contagion. The appropriateness of using this procedure as a measure to increase milk yield, and its possible influence on the public opinion, should also be weighted into the equation when considering its use.

#### ***2.4.2 Visual Enrichment***

701 The visual system of cattle is very sensitive to motion and contrasts of light and dark (Grandin  
702 2000). The lateral eye position enables cattle to constantly scan the horizon (up to 330 degrees)  
703 for predators, and facilitates other activities, such as sexual (mounting) behaviour (Grandin 2000,  
704 Tucker 2009). Visual enrichment, as opposed to auditory and olfactory enrichment, can be  
705 simply avoided by the animals when found to be aversive (i.e. by closing their eyelid, or  
706 diverting their gaze away), and consequently, controlled (Wells, 2009). However, to the best of  
707 our knowledge, very little work has been carried out on the effect of visual enrichment on cattle.  
708 Haskell et al. (2013) found no significant effect of access to a view of surrounding fields and  
709 farm tracks on the motivation of cows to occupy a loafing area. The authors' interpretation was  
710 that a view of the surroundings has little motivational value for cows. However, since the effect  
711 of visual access to the surroundings was not assessed during feeding time (i.e. food was placed  
712 inside the cowshed), when lying down (i.e. loafing area had concrete flooring which is less  
713 favored by cows) or during the evening/night (i.e. observations in this study were conducted only  
714 during day light hours), further investigation on this potential enrichment is needed. An  
715 additional factor that should be controlled for when assessing the effect of visual surroundings on  
716 dairy cattle is the confounding effect of previous experience (discussed above under physical and  
717 cognitive enrichment, Tucker et al., 2003; Manteuffel et al., 2009). Piller et al. (1999) studied the  
718 effect of mirror-image exposure on heart rate and movement of isolated heifers. The presence of  
719 mirrors was associated with reduced heart rate and movement, and had a greater calming effect  
720 when placed directly in front of the animal, compared to a reflected side-view. In accordance  
721 with these results, Coulon et al. (2011) found that heifers were more attracted to images of  
722 familiar conspecifics (i.e. chosen first, explored more, and given more attention) compared to  
723 images of unfamiliar conspecifics. If images of conspecifics are indeed treated as representations

of the real individuals, they might prove to be beneficial for lowering stress levels during husbandry/medical procedures that require isolation. In a study measuring salivary cortisol and chromogranin A concentrations from three Japanese Black cows, a picture of a companion, compared to a blank picture, tended to decrease, but not significantly, stress response measured after 30 minutes following the start of the isolation. However, two hours later, the opposite effect was observed for salivary cortisol levels (Ninomiya and Sato, 2011). One possible explanation for the time-limited effect of presenting a picture of a companion on lowering stress levels is that frustration might build up over time, because of the inability to eventually engage in contact with the conspecifics (Wells, 2009). More research is needed in order to gain a better understanding of the short term and long term beneficial effects of this tool in lowering stress. It is important to point out that reducing stress responses in cattle can also be achieved by eliminating visual input rather than enriching it. Mitchell et al. (2004) found that preventing beef cattle from seeing by blindfolding them, reduced the amount of struggle and tended to lower heart rate during restraint. We encourage research aimed at a better understanding of the “calming effect” induced by blindfolding and a systematic comparison between these two approaches (i.e. mirror vs. blindfolding).

### ***2.4.3 Olfactory Enrichment***

Cattle, compared to humans, have a very sensitive sense of smell. Using their vomeronasal organ, cattle can detect pheromones indicating the reproductive state of their conspecifics or their stress state (via their urine; Terlouw et al., 1998). To our knowledge, there is only one study that assessed the suitability of olfactory enrichment in cattle. Wilson et al. (2002) compared two tactile enrichment devices (e.g. a scratching/rubbing walkway and a movable scratching/rubbing

device), two olfactory enriching devices (e.g. a milk-scent releasing device and a lavender-scent releasing device) and a control non-scented device. Their results show that tactile enriching devices were used for a longer duration, more frequently and by a higher percentage of cows compared to the scent devices. Moreover, the initial preference that cows displayed towards the milk and lavender-scent devices over to the non-scented control device decreased over the course of the experiment. The authors concluded that scratching/rubbing devices are better candidates for environmental enrichment than scent devices, as the interest that cows display towards them does not fade over time. A different approach that holds higher biological relevance, and that, to our knowledge, has not been tested yet, is to enrich the scent of cows food ration, either with artificial or natural smell. One possible constraint associated with this idea is the reluctance of ruminants to eat novel food (known as food neophobia; Launchbaugh, 1995), and the possible association of the new smells with new tastes. We will further discuss this limitation in the nutritional enrichment section.

#### ***2.4.4 Tactile Enrichment***

On pasture, cattle use trees and other inanimate objects to groom themselves. The use of trees for rubbing seems to originate from a different need than grooming behaviour. When trees are available, cattle will use them for scratching different body regions, without spending less time self-grooming and allogrooming than when trees are absent (Kohari et al., 2007). Inside the dairy farm, lacking any trees, dairy cows rub their body and mainly their head and neck on metal gates, fences and water troughs (DeVries et al., 2007). A useful tactile enrichment device that can be placed on the farm and ease grooming behaviour of cows is an automated brush. Automatic brushes allow cattle to groom themselves, particularly in body regions that they find hard to

reach otherwise (DeVries et al., 2007), and were shown to be preferred over fixed brushes (Gutmann 2010). In addition to providing the cows with the opportunity to engage in scratching/rubbing behaviour, the brushes are also associated with better body cleanliness and in some situations, improved milk yield (second lactation, Schukken and Young, 2009). Once a brush is installed on the farm, calves, lactating dairy cows, dry cows and breeding bulls (both young and mature), will use it on a daily basis (Georg and Totschek, 2001, Georg et al., 2007; DeVries et al., 2007; Hoyer 2013, Newby et al., 2013; Mandel et al., 2013). The frequency of brush usage varies between cows and housing systems, and was found to be considerably higher (4-7 fold) in studies where cows/bulls were housed individually (Hoyer 2013; Newby et al., 2013), compared to studies where one brush was present for a group of cows (Georg and Totschek, 2001, Georg et al., 2007; DeVries et al., 2007; Mandel et al., 2013). The high usage rates by individually housed cows can be attributed to reduced competition over this resource (Val-Laillet et al., 2008), boredom (Hoyer 2013) or simply the lack of allogrooming possibilities. Another factor that plays an important role in brush utilization is the location of the brush and its distance from the food bunk (Mandel et al., 2013). In addition to its hedonistic character and high utilization rates, brush usage can serve as an interesting tool to measure and identify stress and morbidity in dairy cows (Mandel et al., 2013). Indeed, in this previous paper, we found reduced brush usage under heat load and following intrusive medical procedures (i.e. stress; Mandel et al., 2013). In accordance with this idea, a recent study showed that brush usage is also reduced among steers infected with Bovine Respiratory disease (BRD) on the day of peak illness (i.e. morbidity; Toaff-Rosenstein et al., 2014).

The need of cows to engage in scratching/rubbing behaviour can also be achieved by brushing them manually. In a recent study, Westerath et al. (2014) examined whether calves judge human brushing as positive. The majority of calves faced with a preference test preferred a compartment with a human that brushed the animals over an empty compartment. The authors mentioned that all calves “leaned against the brush” and “stretched the neck while being brushed”, possibly indicating some kind of perceived pleasure when being brushed. However, the authors emphasize the need to habituate the animals to being brushed, in order for the positive impact of brushing to exceed the fear of novelty or the fear of a contact with humans (as opposed to very rapid adaptation of cows to automated brushes, DeVries et al., 2007). Accordingly, Westerath et al., (2014) also mention that some of the animals used in their study never habituated to being brushed by the experimenter, implying that this action is not perceived to be positive by all individuals. Hanging manila ropes (i.e. fiber ropes originally used for pathogen sampling) in the yard may also serve as a tactile enrichment device (Stanford et al., 2009). Following a short adaptation period, cattle will lick and nibble the rope on a daily basis (Stanford et al., 2009). Calves and heifers interact more with the rope than mature cows (Stanford et al., 2005; Stanford et al., 2009). However, to our knowledge, the nature of the motivation to engage in this behaviour was not investigated, and its origin should be assessed to control for nutritional deficits/stereotypic origin.

## ***2.5 Nutritional Enrichment***

Nutritional enrichment can involve either presenting varied or novel food types, or changing the method of food delivery (Bloomsmith et al., 1991).

**Calves.** Calves start to suckle milk from their dam within hours of birth (domestic settings: Edwards and Broom, 1982; non-domestic settings: Tucker, 2009). The motivation to suckle at this age is very high, and the frequency of milk intake in dairy calves ranges between 8-12 times a day (dairy calves, Jensen, 2003). Calves kept with their dam can suckle at will, and tend to have frequent suckling bouts during which they ingest small amounts of milk at a time (Kalber and Barth, 2014). In the following weeks, the frequency of milk intake will gradually decrease to 3-4 times a day (Jensen, 2003), allowing plant based diet to slowly build up. Natural weaning occurs between 6 months and one year (Newberry and Swanson, 2008). Allowing calves to suckle directly from the teats of their dam/foster cow (i.e. either in a full or restrict contact dam rearing system), has been associated with higher growth rates and reduced cross-sucking behaviour (Roth et al., 2009). However, the higher growth rate associated with dam-rearing could be confounded by a higher milk intake (of natural milk) compared to calves provided with milk replacer (Kalber and Barth, 2014).

In most intensive farming systems, calves are raised separately from their dam and milk is typically provided via a teat feeder or in a bucket. When fed from a bucket, calves consume a large amount of milk in a very short time (i.e. 2.5 liters in about one minute, compared to 8-12 minutes for an average suckling bout from the dam's udder, Loberg and Lidfors, 2001). Feeding calves through an open bucket (i.e. without a nipple) is cheap and easy to maintain, but does not provide calves with the possibility to perform suckling behaviour. Preventing calves from performing suckling behaviour may result in its redirection towards pen objects or peers (i.e. cross-sucking). Although the origin of cross-sucking behaviour is thought to be influenced by multiple factors, such as milk allowance and age of weaning, recent findings show that cross-

sucking may reflect individual differences or be the result of habit formation (de Passillé et al., 2011).

The traditional method for treating cross-suckling behaviour is to install a pronged nose-ring or halter on the nose of the suckling calf. Spikes attached to the nose ring/halter provoke avoidance behaviour among group members, and limit their ability to suckle each other. The method was criticized for targeting the signs rather than the cause (Keil et al., 2000). An alternative treatment that deals with the cause itself and gives calves the opportunity to perform suckling behaviour is to place a rubber nipple in the pen. de Passillé and Caza (1997) showed that the occurrence of cross-suckling behaviour is reduced by more than 75% when calves are presented with dry rubber nipples following milk intake from a bucket. However, although the rate of cross-suckling was reduced using this method, it still involves the performance of a functionless behaviour (i.e. sucking a dry rubber teat), which may be referred to as stereotypic (de Passillé et al., 2011). A more promising solution is to provide calves with nutritive feeding nipples (i.e. where milk is being provided through the rubber nipple), which reduces both cross-suckling behaviour (de Passillé, 2001; Jensen and Weary, 2013), and the time spent in non-nutritive suckling (i.e. when milk is finished and the nipples are dry; de Passillé et al., 2011). The nutritive rubber nipple can be connected to a bucket (i.e. teat-bucket) or to a feeder (i.e. teat-feeder), and be used to provide calves with several milk meals a day. Since calves, when suckling from their mothers, tend to switch quarters once milk flow begins to decline (in contrast to piglets which use the same teat; Haley et al., 1998), placing shoulder barriers between feeding stations is encouraged, in order to prevent competition between calves (Jensen et al., 2008). Enriching the environment of a teat-



feeder by adding a post-feeding area with non-nutritive rubber teats and a net filled with a hay bale, can also further reduce cross-sucking behaviour (Ude et al., 2011).

**Cows.** On pasture, dairy cows graze between 6-12 hrs a day, depending on nutrient availability, ingestion speed and competition over food resources (Coffey et al., 1992). However, when kept indoors, feeding time is reduced to 4 hrs a day on average (Gomez and Cook, 2010). This change in feeding time can be explained, at least to some extent, by the highly predictable food location and the easily consumed form in which food is being dispensed in indoor conditions (e.g. fenceline feeding of total mixed ration (TMR); Newberry, 1993; Gomez and Cook, 2010). This form of dispensing food is however associated with increased incidence of agonistic interactions, which is usually not the case on pasture, when the animals are spaced out from each other (Miller and Wood-Gush, 1991). Providing cows with a larger feeding space (1.0 m per cow compared to 0.5 m) resulted in at least a 60% increase in space between animals and a 57% reduction in aggressive interactions while feeding (DeVries et al., 2004). This change was associated with increased feeding activity throughout the day, especially during the 90 min after providing fresh feed (an increase of 24%). More importantly, the increase in feeding activity was particularly evident for subordinate cows. When providing extra feeding space per cow is not possible, placing feed barriers (i.e. headlocks) can serve as an intermediate solution that decreases displacements at the feeding bunk by more than 20% (Endres et al., 2005).

Although keeping cows indoors seems to affect the "naturalness" of their feeding behaviour (i.e. the extent to which a behaviour resembles that performed in more extensive or natural environment), the functionality and adaptiveness of indoor feeding techniques should be

emphasized (Newberry, 1995). The demand for high milk yield in intense dairy farming systems places high metabolic demand on dairy cows (Butler and Smith, 1989; Rauw et al., (1998), Oltenacu and Algers (2005), or Veerkamp (2009). Providing cows with easier access to food (i.e. in the form of TMR) reduces the time they need to spend standing in order to feed, and may thus be adaptive in order for cows to cope with the energetic demand placed on them. However, the functionality and adaptiveness of this feeding method for heifers is not clear, as they do not share the same metabolic demands as lactating dairy cows, and are consequently left to occupy longer periods of time with a limited range of possible behavioural patterns. Providing part of the daily food ration through a food net (or a device that forces the animal to engage in food collection - such as pulling the hay from a bundle rather than just picking it up from the floor) may prolong the duration of feeding behaviour and serve as a measure to increase the "naturalness" of calf feeding behaviour. Providing essential resources (such as food) both in their regular form (i.e. feed bunks) and as a reward (i.e. food net), can enrich cows' feeding experience through contrafreeloading (i.e. the phenomenon where an animal will work for a reward in the presence of the same reward available freely; reviewed by Inglis et al., 1997) and may contribute to their welfare. We encourage further investigation of this idea as an enrichment method for heifers.

Offering animals a wider selection of food types was also suggested to be a potential source of enrichment (Newberry 1995). Bioactive forages for example, are offered to combat gastrointestinal parasites and serve as an alternative to anthelmintic drugs (Hutchings et al., 2003). However, when considering the practicality of this enrichment method, one should also take into account the effect of neophobia towards unfamiliar food (Westerath et al., 2014, although see Costa et al., 2014 for reduced neophobia in calves raised in complex social groups

including both calves and cows). Familiarizing the cows with varying types of food (of which some is regarded to be more rewarding and some less, e.g. concentrate in calves, Westerath et al., 2014) may prove to be beneficial in decreasing neophobia, but may have an impact on the bacterial diversity in the rumen, which plays a major role in the productivity and health of cows (Callaway et al., 2010). In addition, food that is considered to be more rewarding (e.g. concentrate in calves, Westerath et al., 2014) may increase competition, and render the food accessible only to more dominant cows. In accordance with this idea, Rioja-Lang et al. (2009) showed that dairy cows trade-off food quality with proximity to a dominant individual in Y-maze choice tests. Therefore, enriching the food of farm animals (using familiar food which is perceived to be more rewarding by the animals) might prove to be more suitable in farms that dispense food individually (via individual recognition systems), or in farms that install proper barriers between feeding stations to avoid agonistic behaviour (Arachchige et al., 2014). A different approach, which to our knowledge has not been tested yet, is to use nutritional enrichment as a measure to detect lack of sufficient nutrients in the diet. Bell and Sly (1983) showed that the hedonic characteristics of nutritional enrichment, such as salt-licks, may be limited to nutrient deficient cattle. However, this idea should be further examined.

### 3. CONCLUSION

Indoor housing of dairy cows and calves is associated with various challenges for the animals, such as the constraint to occupy long periods of time with a limited range of behaviour patterns, as well as maintenance in abnormal social groups (Hughes and Duncan, 1988, Morgan and Tromborg, 2007). Here, we have reviewed environmental enrichment methods that are aimed at assisting cattle to cope better with stressors in their environment, prevent frustration and increase

the fulfillment of behavioural needs. As animal welfare is considered not only as the absence of stress and harm, but also as the promotion of better affective conditions (Boissy et al. 2007), the implementation of cognitive enrichment that can lead to positive emotions was also discussed. Several of the basic behavioural requirements described here (e.g. social enrichment in the form of contact with conspecifics) are considered as minimal requirements for raising standards for gregarious animals, and in some countries are enshrined under legislative acts (European Union Council Directive 2008/119/EC, Article 3). However, in many other countries, even these basic requirements are not met (Fraser et al., 2013). Enrichment methods aimed at fulfilling other important behavioural needs (e.g. providing cows with a secluded area to calve, or feeding calves with milk through a nipple), have the potential to advance the welfare of dairy cows and calves, but are not yet statutory. Finally, the contribution of enrichment methods that are less biologically relevant (reviewed here under olfactory and auditory enrichment, e.g. classical music, lavender smell) to the wellbeing of cattle is, however, less clear. Newberry (1995) argued that "enrichment attempts will fail if the environmental modifications have little functional significance to the animals, are not sufficiently focused to meet a specific goal, or are based on an incorrect hypothesis regarding the causation and mechanisms underlying a problem". Once these criteria are met, other factors, such as the accessibility of the enrichment device to cows from different hierarchical rankings (and competition over resources), as well as the difficulty of cleaning and disinfecting enclosures and other enrichment materials, should be taken into consideration when assessing the practicality of a specific enrichment device. The benefits of enriching the environment of dairy cows and calves, in terms of improving the physical and mental state of the animals, and in some cases increased productivity, should be weighted in the expected costs of integrating these methods into commercial farms, and may consequently play

an important part in increasing the motivation of farmers to adopt these methods. Several of the methods described here, will, with time, be integrated to the minimum raising standards of cattle, whereas others will remain under the scope of enrichment, and continue to pull the field of animal welfare forward, allowing us to deepen our understanding of farm animals and their needs inside the industry. As zero grazing systems gain popularity around the world, more research will be needed to assess their impact on the wellbeing of the animals, and the development of enrichment methods which are better adapted to cows and calves housed indoors will be required.

#### ACKNOWLEDGEMENTS

We would like to thank E. Briefer, M. Wenker and three anonymous reviewers for helpful comments on the manuscript. R. M. is funded by the Universities Federation for Animal Welfare and by the Harry and Sylvia Hoffman Leadership and Responsibility Program at the Hebrew University, Israel.

REFERENCES

- Algers, B., Bertoni, G., Broom, D., Hartung, J., Lidfors, L., Metz, J., Munksgaard, L., Pina, T.N., Oltenacu, P., Rehage, J., and Rushen, J. 2009. Scientific report on the effects of farming systems on dairy cow welfare and disease<sup>1</sup>. Report of the Panel on Animal Health and Welfare (Question No EFSA-Q-2006-113). doi:10.2903/j.efsa.2009.1143r.
- Almeida, P.E., Weber, P.S.D., Burton, J.L., and Zanella, A.J. 2008. Depressed DHEA and increased sickness response behaviors in lame dairy cows with inflammatory foot lesions. *Domest. Anim. Endocrinol.*, 34: 89–99.
- Arachchige, A. H., Fisher, A. D., Wales, W. J., Auldist, M. J., Hannah, M. C., & Jongman, E. C. 2014. Space allowance and barriers influence cow competition for mixed rations fed on a feed-pad between bouts of grazing. *J. Dairy Sci.*, 97, 3578-3588.
- Arnold, N. A., Ng, K. T., Jongman, E. C., and Hemsworth, P. H. 2007. The behavioural and physiological responses of dairy heifers to tape-recorded milking facility noise with and without a pre-treatment adaptation phase. *Appl. Anim. Behav. Sci.*, 106, 13-25.
- Arnold, N. A., Ng, K. T., Jongman, E. C., and Hemsworth, P. H. 2008. Avoidance of tape-recorded milking facility noise by dairy heifers in a Y maze choice task. *Appl. Anim. Behav. Sci.*, 109, 201-210.

- 991 Bach, A. 2012. Ruminant Nutrition Symposium: Optimizing Performance of the Offspring:  
992 Nourishing and managing the dam and postnatal calf for optimal lactation, reproduction, and  
993 immunity. *J. Anim. Sci.*, 90, 1835-1845.
- 994
- 995 Bailey, D. W., Rittenhouse, L. R., Hart, R. H., Swift, D. M., and Richards, R. W. 1989a.  
996 Association of relative food availabilities and locations by cattle. *J. Range. Manage.*, 42, 480-  
997 482.
- 998
- 999 Bailey, D. W., Rittenhouse, L. R., Hart, R. H., and Richards, R. W. 1989b. Characteristics of  
1000 spatial memory in cattle. *Appl. Anim. Behav. Sci.*, 23, 331-340.
- 1001
- 1002 Barkema, H. W., von Keyserlingk, M. A. G., Kastelic, J. P., Lam, T. J. G. M., Luby, C., Roy, J.  
1003 P., LeBlanc, S.J. Keefe, J.P and Kelton, D. F. 2015. Invited review: Changes in the dairy  
1004 industry affecting dairy cattle health and welfare. *J. dairy Sci.* doi:10.3168/jds.2015-9377.
- 1005
- 1006 Bassett, L., and Buchanan-Smith, H. M. 2007. Effects of predictability on the welfare of captive  
1007 animals. *Appl. Anim. Behav. Sci.*, 102, 223-245.
- 1008
- 1009 Baumans, V., and Van Loo, P. L. P. 2013. How to improve housing conditions of laboratory  
1010 animals: The possibilities of environmental refinement. *Vet. J.*, 195, 24-32.
- 1011
- 1012 Bell, F. R., and Sly, J. 1983. The olfactory detection of sodium and lithium salts by sodium  
1013 deficient cattle. *Physiol. Behav.*, 31, 307-312.

1014

1015 Bloomsmith, M. A., L. Y. Brent, and S. J. Schapiro. 1991. Guidelines for developing and  
1016 managing an environmental enrichment program for nonhuman primates. *Lab. Anim. Sci.*,  
1017 41:372–377.

1018

1019 Bøe, K. E., and Færevik, G. 2003. Grouping and social preferences in calves, heifers and cows.  
1020 *Appl. Anim. Behav. Sci.*, 80, 175-190.

1021

1022 Boissy, A., Manteuffel, G., Jensen, M. B., Moe, R. O., Spruijt, B., Keeling, L. J., Winckler, C.,  
1023 Forkman, B., Dimitrov, I., Langbein, J., Bakken, M., Veissier, I., and Aubert, A. 2007.  
1024 Assessment of positive emotions in animals to improve their welfare. *Physiol. Behav.*, 92, 375-  
1025 397.

1026

1027 Borderas, T.F., de Passillé, A.M, and Rushen, J. 2008. Behavior of dairy calves after a low dose  
1028 of bacterial endotoxin. *Anim. Sci. J.*, 86: 2920-2927.

1029

1030 Boyle, A. R., Ferris, C. P., and O’Connell, N. E. 2012. Are there benefits in introducing dairy  
1031 heifers to the main dairy herd in the evening rather than the morning?. *J. Dairy Sci.*, 95, 3650-  
1032 3661.

1033

1034 Breuer, K., Hemsworth, P. H., Barnett, J. L., Matthews, L. R., and Coleman, G. J. 2000.  
1035 Behavioural response to humans and the productivity of commercial dairy cows. *Appl. Anim.*  
1036 *Behav. Sci.*, 66, 273-288.



- 1037
- 1038 Briefer, E. F. 2012. Vocal expression of emotions in mammals: mechanisms of production and  
 1039 evidence. *J. Zool.*, 288, 1-20.
- 1040
- 1041 Broom, D. M. 2010. Cognitive ability and awareness in domestic animals and decisions about  
 1042 obligations to animals. *Appl. Anim. Behav. Sci.*, 126, 1-11.
- 1043
- 1044 Broom, D.M. and Leaver, J.D. 1978. Effects of group-rearing or partial isolation on later social  
 1045 behaviour of calves. *Anim. Behav.*, 26, 1255–1263.
- 1046
- 1047 Bulens, A., Van Beirendonck, S., Van Thielen, J., and Driessen, B. 2014. The effect of  
 1048 environmental enrichment on the behaviour of beef calves. In *Proceedings of the 6th*  
 1049 *International Conference on the Assessment of Animal Welfare at Farm and Group Level* (pp.  
 1050 135-135).
- 1051
- 1052 Butler, W. R., and Smith, R. D. 1989. Interrelationships between energy balance and postpartum  
 1053 reproductive function in dairy cattle. *J. Dairy Sci.*, 72, 767-783.
- 1054
- 1055 Callaway, T. R., Dowd, S. E., Edrington, T. S., Anderson, R. C., Krueger, N., Bauer, N.,  
 1056 Kononoff, P.J., and Nisbet, D. J. 2010. Evaluation of bacterial diversity in the rumen and feces of  
 1057 cattle fed different levels of dried distillers grains plus solubles using bacterial tag-encoded FLX  
 1058 amplicon pyrosequencing. *Anim. Sci. J.*, 88, 3977-3983.
- 1059

Carlstead, K.C., Shepherdson, D.J., 2000. Alleviating stress in zoo animals with environmental enrichment. Pages 337–354 In *The Biology of Animal Stress: Basic Principles and Implications for Animal Welfare*. Moberg, G.P., Mench, Joy, A. (Eds.). CAB International Publishing, Oxon, NY.

Castro, I. M., Gygax, L., Wechsler, B., and Hauser, R. 2011. Increasing the interval between winter outdoor exercise aggravates agonistic interactions in Hérens cows kept in tie-stalls. *Appl. Anim. Behav. Sci.*, 129, 59-66.

Canadian Dairy information center. Dairy barns by type in Canada. Accessed Sept. 9, 2015. [http://www.dairyinfo.gc.ca/index\\_e.php?s1=dff-fcilands2=farm-fermeands3=db-el](http://www.dairyinfo.gc.ca/index_e.php?s1=dff-fcilands2=farm-fermeands3=db-el).

CBS 2015 (Dutch Central Bureau for Statistics). Access to pasture in dairy cattle; Pasture area. Accessed Sept. 02, 2015. <http://statline.cbs.nl/StatWeb/publication/?VW=TandDM=SLnlandPA=70736nedandLA=nl>; Last seen 02/09/2015.

Cobb, C. J., Obeidat, B. S., Sellers, M. D., Pepper-Yowell, A. R., Hanson, D. L., and Ballou, M. A. 2014. Improved performance and heightened neutrophil responses during the neonatal and weaning periods among outdoor group-housed Holstein calves. *J. Dairy Sci.*, 97, 930-939.

- 1081 Coffey, K. P., Moyer, J. L., Brazle, F. K., and Lomas, L. W. 1992. Amount and diurnal  
1082 distribution of grazing time by stocker cattle under different tall fescue management strategies.  
1083 *Appl. Anim. Behav. Sci.*, 33, 121-135.  
1084
- 1085 Collins, M. T., Eggleston, V., and Manning, E. J. B. 2010. Successful control of Johne's disease  
1086 in nine dairy herds: results of a six-year field trial. *J. Dairy Sci.*, 93, 1638-1643.  
1087
- 1088 Cooper, M. D., Arney, D. R., and Phillips, C. J. 2008. The effect of temporary deprivation of  
1089 lying and feeding on the behaviour and production of lactating dairy cows. *Animal*, 2, 275-283.  
1090
- 1091 Costa, J. H. C., Daros, R. R., von Keyserlingk, M. A. G., and Weary, D. M. 2014. Complex  
1092 social housing reduces food neophobia in dairy calves. *J. Dairy Sci.*, 97, 7804-7810.  
1093
- 1094 Coulon, M., Baudoin, C., Heyman, Y., and Deputte, B. L. 2011. Cattle discriminate between  
1095 familiar and unfamiliar conspecifics by using only head visual cues. *Anim. Cogn.*, 14, 279-290.  
1096
- 1097 Daros, R. R., Costa, J. H., von Keyserlingk, M. A., Hötzel, M. J., and Weary, D. M. 2014.  
1098 Separation from the dam causes negative judgement bias in dairy calves. *PLoS ONE* 9: e98429.  
1099 doi: 10.1371/journal.pone.0098429. pmid:24848635.  
1100
- 1101 Dawkins, M. S. 1983. Battery hens name their price: consumer demand theory and the  
1102 measurement of ethological 'needs'. *Anim. Behav.*, 31, 1195-1205.  
1103

- 1104 de Passillé, A. M. 2001. Sucking motivation and related problems in calves. *Appl. Anim. Behav.*  
 1105 *Sci.*, 72, 175-187.  
 1106
- 1107 de Passillé, A. M. B., and Caza, N. 1997. Cross-sucking by calves occurs after meals and is  
 1108 reduced when calves suck a dry teat. *J. Dairy Sci.*, 80, 229.  
 1109
- 1110 de Passillé, A. M., Borderas, F., and Rushen, J. 2011. Cross-sucking by dairy calves may become  
 1111 a habit or reflect characteristics of individual calves more than milk allowance or weaning. *Appl.*  
 1112 *Anim. Behav. Sci.*, 133, 137-143.  
 1113
- 1114 de Passillé, A. M., Rushen, J., Ladewig, J., and Petherick, C. 1996. Dairy calves' discrimination  
 1115 of people based on previous handling. *Anim. Sci. J.*, 74, 969-974.  
 1116
- 1117 de Paula Vieira, A., de Passillé, A. M., and Weary, D.M. 2012. Effects of the early social  
 1118 environment on behavioral responses of dairy calves to novel events. *J. Dairy Sci.*, 95, 5149–  
 1119 5155.  
 1120
- 1121 de Paula Vieira, A., von Keyserlingk, M. A. G., and Weary, D. M. 2010. Effects of pair versus  
 1122 single housing on performance and behavior of dairy calves before and after weaning from milk.  
 1123 *J. Dairy Sci.*, 93, 3079-3085.  
 1124
- 1125 DeVries, T. J., Vankova, M., Veira, D. M., and von Keyserlingk, M. A. G. 2007. Usage of  
 1126 mechanical brushes by lactating dairy cows. *J. Dairy Sci.*, 90, 2241-2245.

1127

1128 DeVries, T. J., von Keyserlingk, M. A. G., and Weary, D. M. 2004. Effect of feeding space on  
 1129 the inter-cow distance, aggression, and feeding behavior of free-stall housed lactating dairy  
 1130 cows. *J. Dairy Sci.*, 87, 1432-1438.

1131

1132 Duve, L. R., and Jensen, M. B. 2011. The level of social contact affects social behaviour in pre-  
 1133 weaned dairy calves. *Appl. Anim. Behav. Sci.*, 135, 34-43.

1134

1135 Duve, L. R., Weary, D. M., Halekoh, U., and Jensen, M. B. 2012. The effects of social contact  
 1136 and milk allowance on responses to handling, play, and social behavior in young dairy calves. *J.*  
 1137 *Dairy Sci.*, 95, 6571-6581.

1138

1139 Edgar, J., Held, S., Paul, E., Pettersson, I., Price, R. I. A., and Nicol, C. 2015. Social buffering in  
 1140 a bird. *Anim. Behav.*, 105, 11-19.

1141

1142 Edwards, S. A., and Broom, D. M. 1982. Behavioural interactions of dairy cows with their  
 1143 newborn calves and the effects of parity. *Anim. Behav.*, 30, 525-535.

1144

1145 Ellingsen, K., Coleman, G. J., Lund, V., and Mejdell, C. M. 2014. Using qualitative behaviour  
 1146 assessment to explore the link between stockperson behaviour and dairy calf behaviour. *Appl.*  
 1147 *Anim. Behav. Sci.*, 153, 10-17.

1148

- 1149 Endres, M. I., DeVries, T. J., von Keyserlingk, M. A. G., and Weary, D. M. 2005. Short  
1150 communication: effect of feed barrier design on the behavior of loose-housed lactating dairy  
1151 cows. *J. Dairy Sci.*, 88, 2377-2380.
- 1152
- 1153 Færevik, G., Jensen, M. B., and Bøe, K. E. 2006. Dairy calves social preferences and the  
1154 significance of a companion animal during separation from the group. *Appl. Anim. Behav. Sci.*,  
1155 99, 205-221.
- 1156
- 1157 Flower, F. C., and Weary, D. M. 2001. Effects of early separation on the dairy cow and calf:: 2.  
1158 Separation at 1 day and 2 weeks after birth. *Appl. Anim. Behav. Sci.*, 70, 275-284.
- 1159
- 1160 Flower, F. C., and Weary, D. M. 2003. The effects of early separation on the dairy cow and calf.  
1161 *Anim. Welf.*, 12, 339-348.
- 1162
- 1163 Fogsgaard, K.K., Røntved, C.M., Sørensen, P., and Herskin, M.S. 2012. Sickness behavior in  
1164 dairy cows during *Escherichia coli* mastitis. *J. Dairy Sci.*, 95, 630–638.
- 1165
- 1166 Fraser, D. 2008. *Understanding Animal Welfare: The Science in Its Cultural Context*. Wiley-  
1167 Blackwell, Oxford, UK.
- 1168
- 1169 Fraser, D., Duncan, I. J., Edwards, S. A., Grandin, T., Gregory, N. G., Guyonnet, V.,  
1170 Hemsworth, P.H., Huertas, S.M., Huzzey, J.M., Mellor, D.J., Mench, J.A., Špinka, M. and

- 1171 Whay, H. R. 2013. General Principles for the welfare of animals in production systems: The  
1172 underlying science and its application. *Vet. J*, 198, 19-27.  
1173
- 1174 Fraser, D., Weary, D. M., Pajor, E. A., and Milligan, B. N. 1997. A scientific conception of  
1175 animal welfare that reflects ethical concerns. *Anim. welf.* 6, 187-205.  
1176
- 1177 Fregonesi, J. A., Veira, D. M., von Keyserlingk, M. A. G., and Weary, D. M. 2007. Effects of  
1178 bedding quality on lying behavior of dairy cows. *J. Dairy Sci.*, 90, 5468-5472.  
1179
- 1180 Gaillard, C., Meagher, R. K., von Keyserlingk, M. A., and Weary, D. M. 2014. Social housing  
1181 improves dairy calves' performance in two cognitive tests. *PloS ONE*, 9, e90205.  
1182
- 1183 Galindo, F., and Broom, D. M. 2002. The effects of lameness on social and individual behavior  
1184 of dairy cows. *Appl. Anim. Behav. Sci.*, 5, 193-201.  
1185
- 1186 Georg, H., and Totschek, K. 2001. Examining an automatic cow brush for dairy cows.  
1187 *Landtechnik*, 56, 260-261.  
1188
- 1189 Georg, H., Jahn-Falk, D., and Brunswick, G.U. 2007. Technology against boredom.  
1190 *Landtechnik*, 62, 166–167.  
1191
- 1192 Gomez, A., and Cook, N. B. 2010. Time budgets of lactating dairy cattle in commercial freestall  
1193 herds. *J. Dairy Sci.*, 93, 5772-5781.

- 1194
- 1195 Grandin, T. 2000. Handling and welfare of livestock in slaughter plants. Pages 409-439 in
- 1196 Grandin (ed) *Livestock Handling and Transport* (2nd). CAB International, Wallingford, Oxon,
- 1197 UK.
- 1198
- 1199 Guillot, F.S. 1981. Susceptibility of Hereford cattle to sheep scab mites after recovery from
- 1200 psoroptic scabies. *J. Econ. Entomol.*, 74, 653–657.
- 1201
- 1202 Gutmann, A. 2010. Verhalten von Milchkühen bei der Nutzung von fixen gegenüber rotierenden
- 1203 Bürsten. Pages 78–81 in the *Proceeding of 24th IGN Conference*, 3-5 June 2010, Ettenhausen,
- 1204 Switzerland.
- 1205
- 1206 Gutmann, A. K., Špinka, M., and Winckler, C. 2015. Long-term familiarity creates preferred
- 1207 social partners in dairy cows. *Appl. Anim. Behav. Sci.*, 169, 1-8.
- 1208
- 1209 Gygax, L., Neisen, G., and Wechsler, B. 2009. Differences between single and paired heifers in
- 1210 residency in functional areas, length of travel path, and area used throughout days 1–6 after
- 1211 integration into a free stall dairy herd. *Appl. Anim. Behav. Sci.*, 120, 49-55.
- 1212
- 1213 Gygax, L., Neisen, G., and Wechsler, B. 2010. Socio-spatial relationships in dairy cows.
- 1214 *Ethology*, 116, 10-23.
- 1215



- 1216 Hagen, K., and Broom, D. M. 2004. Emotional reactions to learning in cattle. *Appl. Anim.*  
 1217 *Behav. Sci.*, 85, 203-213.  
 1218
- 1219 Haley, D. B., Rushen, J., Duncan, I. J., Widowski, T. M., and de Passillé, A. M. 1998. Butting by  
 1220 calves, *Bos taurus*, and rate of milk flow. *Anim. Behav.*, 56, 1545-1551.  
 1221
- 1222 Haskell, M. J., Rennie, L. J., Bowell, V. A., Bell, M. J., and Lawrence, A. B. 2006. Housing  
 1223 system, milk production, and zero-grazing effects on lameness and leg injury in dairy cows. *J.*  
 1224 *Dairy Sci.*, 89, 4259-4266.  
 1225
- 1226 Haskell, M. J., Masłowska, K., Bell, D. J., Roberts, D. J., and Langford, F. M. 2013. The effect  
 1227 of a view to the surroundings and microclimate variables on use of a loafing area in housed dairy  
 1228 cattle. *Appl. Anim. Behav. Sci.*, 147, 28-33.  
 1229
- 1230 Heffner, R. S., and Heffner, H. E. 1983. Hearing in large mammals: Horses (*Equus caballus*) and  
 1231 cattle (*Bos taurus*). *Behav. Neurosci.*, 97, 299-309.  
 1232
- 1233 Held, S. D., and Spinka, M. 2011. Animal play and animal welfare. *Anim. Behav.*, 81, 891-899.  
 1234
- 1235 Hemsworth, P. H. 2003. Human–animal interactions in livestock production. *Appl. Anim.*  
 1236 *Behav. Sci.*, 81, 185-198.  
 1237

1238 Hemsworth, P. H., Coleman, G. J., Barnett, J. L., and Borg, S. 2000. Relationships between  
1239 human-animal interactions and productivity of commercial dairy cows. *Anim. Sci. J.*, 78, 2821-  
1240 2831.

1241

1242 Hepola, H., Hänninen, L., Pursiainen, P., Tuure, V. M., Syrjälä-Qvist, L., Pyykkönen, M., and  
1243 Saloniemi, H. 2006. Feed intake and oral behaviour of dairy calves housed individually or in  
1244 groups in warm or cold buildings. *Livest. Sci.*, 105, 94-104.

1245

1246 Hillmann, E., Roth, B. A., Johns, J., Waiblinger, S., and Barth, K. 2012. Dam-associated rearing  
1247 as animal friendly alternative to artificial rearing in dairy cattle. *Landbauforsch SH*, 362, 181-  
1248 183.

1249

1250 Holm, L., Jensen, M. B., and Jeppesen, L. L. 2002. Calves' motivation for access to two different  
1251 types of social contact measured by operant conditioning. *Appl. Anim. Behav. Sci.*, 79, 175-194.

1252

1253 Hoyer, B. H. 2013. 'Environmental Enrichment'-Strategies to Improve the Housing Conditions of  
1254 Breeding Bulls. Impact on Time Budget, Physical Activity, Rumination, Sexual Behavior and  
1255 Semen Quality. PhD thesis, University of Veterinary Medicine, Hannover, Germany.

1256

1257 Hudson, S. J., and Mullord, M. M. 1977. Investigations of maternal bonding in dairy cattle.  
1258 *Appl. Anim. Ethol.*, 3, 271-276.

1259

- 1260 Hughes, B. O., and Duncan, I. J. H. 1988. The notion of ethological 'need', models of motivation  
1261 and animal welfare. *Anim. Behav.*, 36, 1696-1707.  
1262
- 1263 Hutchings, M. R., Athanasiadou, S., Kyriazakis, I., and J Gordon, I. 2003. Can animals use  
1264 foraging behaviour to combat parasites?. *Proc. Nutr. Soc.*, 62, 361-370.  
1265
- 1266 Ikeda, Y., and Ishii, Y. 2008. Recognition of two psychological conditions of a single cow by her  
1267 voice. *Comput. Electron. Agric.*, 62, 67-72.  
1268
- 1269 Inglis, I. R., Forkman, B., and Lazarus, J. 1997. Free food or earned food? A review and fuzzy  
1270 model of contrafreeloading. *Anim. Behav.*, 53, 1171-1191.  
1271
- 1272 Ishiwata, T., Uetake, K., Kilgour, R. J., Eguchi, Y., and Tanaka, T. 2007. Oral behaviors of beef  
1273 steers in pen and pasture environments. *Appl. Anim. Behav. Sci.*, 10, 185-192.  
1274
- 1275 Jensen, M. B. 2001. A note on the effect of isolation during testing and length of previous  
1276 confinement on locomotor behaviour during open-field test in dairy calves. *Appl. Anim. Behav.*  
1277 *Sci.*, 70, 309-315.  
1278
- 1279 Jensen, M.B. 2003. The effects of feeding method, milk allowance and social factors on milk  
1280 feeding behaviour and cross-sucking in group housed dairy calves. *Appl. Anim. Behav. Sci.* 80:  
1281 191-206.  
1282

- 1283 Jensen, M. B., Duve, L. R., and Weary, D. M. 2015a. Pair housing and enhanced milk allowance  
1284 increase play behavior and improve performance in dairy calves. *J. Dairy Sci.*, 98, 2568-2575  
1285
- 1286 Jensen, M. B., Herskin, M. S., Thomsen, P. T., Forkman, B., and Houe, H. 2015b. Preferences of  
1287 lame cows for type of surface and level of social contact in hospital pens. *J. Dairy Sci.*, 98, 1-8.  
1288
- 1289 Jensen, M. B., and Larsen, L. E. 2014. Effects of level of social contact on dairy calf behavior  
1290 and health. *J. Dairy Sci.*, 97, 5035-5044.  
1291
- 1292 Jensen, M. B., de Passillé, A. M., von Keyserlingk, M. A. G., and Rushen, J. 2008. A barrier can  
1293 reduce competition over teats in pair-housed milk-fed calves. *J. Dairy Sci.*, 91, 1607-1613.  
1294
- 1295 Jensen, M. B., Vestergaard, K. S., and Krohn, C. C. 1998. Play behaviour in dairy calves kept in  
1296 pens: the effect of social contact and space allowance. *Appl. Anim. Behav. Sci.*, 56, 97-108.  
1297
- 1298 Jensen, M. B., Vestergaard, K. S., Krohn, C. C., and Munksgaard, L. 1997. Effect of single  
1299 versus group housing and space allowance on responses of calves during open-field tests. *Appl.*  
1300 *Anim. Behav. Sci.*, 54, 109-121.  
1301
- 1302 Jensen, M.B and Weary, D. 2013. Group Housing and Milk Feeding of Dairy Calves. Pages 179  
1303 - 189 in the proceeding of the 25th Western Canadian Dairy Seminar, *Advances in Dairy*  
1304 *Technology*, March 5-8. University of Alberta, Edmonton. Canada.  
1305

- 1306 Jensen, P., Recen, B., and Ekesbo, I. 1988. Preference of loose housed dairy cows for two  
1307 different cubicle floor coverings. *Swed. J. Agric. Res.*, 18, 141–146.  
1308
- 1309 Johns, J., Patt, A., and Hillmann, E. 2015. Do Bells Affect Behaviour and Heart Rate Variability  
1310 in Grazing Dairy Cows?. *PloS one*, 10, e0131632.  
1311
- 1312 Johnsen, J. F., Marie de Passille, A., Mejdell, C. M., Bøe, K. E., Grøndahl, A. M., Beaver, A.,  
1313 Rushen, J. and Weary, D. M. 2015. The effect of nursing on the cow-calf bond. *Appl. Anim.*  
1314 *Behav. Sci.*, 163, 50-57.  
1315
- 1316 Kalber, T. and Barth, K. 2014. Practical implications of suckling systems for dairy calves in  
1317 organic production systems – a review. *Appl. Agric. Forestry. Res.*, 64, 45-58.  
1318
- 1319 Keil, N. M., Audigé, L., and Langhans, W. 2000. Factors associated with intersucking in Swiss  
1320 dairy heifers. *Prev. Vet. Med.*, 45, 305-323.  
1321
- 1322 Keil, N. M., Wiederkehr, T. U., Friedli, K., and Wechsler, B. 2006. Effects of frequency and  
1323 duration of outdoor exercise on the prevalence of hock lesions in tied Swiss dairy cows. *Prev.*  
1324 *Vet. Med.*, 74, 142-153.  
1325
- 1326 Kikusui, T., Winslow, J. T., and Mori, Y. 2006. Social buffering: relief from stress and anxiety.  
1327 *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, 361, 2215-2228.  
1328

- 1329 Kilgour, R. and Dalton, C. 1984. *Livestock Behaviour: A Practical Guide*. Granada Publishing  
 1330 Ltd: London, UK.  
 1331
- 1332 Kıyıcı, J. M., Koçyigit, R., and Tüzemen, N. 2013. The effect of classical music on milk  
 1333 production, milk components and milking characteristics of Holstein Friesian. *Journal of*  
 1334 *Tekirdag Agricultural Faculty*, 10, 74-81.  
 1335
- 1336 Kohari, D., Kosako, T., Fukasawa, M., and Tsukada, H. 2007. Effect of environmental  
 1337 enrichment by providing trees as rubbing objects in grassland: Grazing cattle need  
 1338 tree-grooming. *Anim. Sci. J.*, 78, 413-416.  
 1339
- 1340 Krohn, C. C. 1994. Behaviour of dairy cows kept in extensive (loose housing/pasture) or  
 1341 intensive (tie stall) environments. III. Grooming, exploration and abnormal behaviour. *Appl.*  
 1342 *Anim. Behav. Sci.*, 42, 73-86.  
 1343
- 1344 Laister, S., Stockinger, B., Regner, A. M., Zenger, K., Knierim, U., and Winckler, C. 2011.  
 1345 Social licking in dairy cattle—Effects on heart rate in performers and receivers. *Appl. Anim.*  
 1346 *Behav. Sci.*, 130, 81-90.  
 1347
- 1348 Langbein, J., Siebert, K., and Nürnberg, G. 2009. On the use of an automated learning device by  
 1349 group-housed dwarf goats: Do goats seek cognitive challenges?. *Appl. Anim. Behav. Sci.*, 120,  
 1350 150-158.  
 1351

- 1352 Launchbaugh, K.L., 1995. Effects of neophobia and aversions of feed intake: Why feedlot cattle  
1353 sometimes refuse to eat nutritious feed. In: Symposium: Intake by Feedlot Cattle. Okla. Agric.  
1354 Exp. Sta., 942, 36-48.  
1355
- 1356 Le Neindre, P. 1989. Influence of rearing conditions and breed on social behaviour and activity  
1357 of cattle in novel environments. Appl. Anim. Behav. Sci., 23, 129-140.  
1358
- 1359 Lensink, B. J., Raussi, S., Boivin, X., Pyykkönen, M., and Veissier, I. 2001. Reactions of calves  
1360 to handling depend on housing condition and previous experience with humans. Appl. Anim.  
1361 Behav. Sci., 70, 187-199  
1362
- 1363 Lidfors, L. M., Moran, D., Jung, J., Jensen, P., and Castren, H. 1994. Behaviour at calving and  
1364 choice of calving place in cattle kept in different environments. Appl. Anim. Behav. Sci., 42, 11-  
1365 28.  
1366
- 1367 Loberg, J., and Lidfors, L. M. 2001. Effect of milkflow rate and presence of floating nipple on  
1368 abnormal sucking between dairy calves. Appl. Anim. Behav. Sci., 72, 189-199.  
1369
- 1370 Loberg, J., Telezhenko, E., Bergsten, C., and Lidfors, L. 2004. Behaviour and claw health in tied  
1371 dairy cows with varying access to exercise in an outdoor paddock. Appl. Anim. Behav. Sci., 89,  
1372 1-16.  
1373

- 1374 Losinger, W. C., and Heinrichs, A. 1997. Management practices associated with high mortality  
1375 among preweaned dairy heifers. *J. Dairy Sci.*, 64, 1-11.  
1376
- 1377 Lürzel, S., Münsch, C., Windschnurer, I., Futschik, A., Palme, R., and Waiblinger, S. 2015. The  
1378 influence of gentle interactions on avoidance distance towards humans, weight gain and  
1379 physiological parameters in group-housed dairy calves. *Appl. Anim. Behav. Sci.* DOI:  
1380 <http://dx.doi.org/10.1016/j.applanim.2015.09.004>  
1381
- 1382 Mandel, R., Whay, H. R., Nicol, C. J., and Klement, E. 2013. The effect of food location, heat  
1383 load, and intrusive medical procedures on brushing activity in dairy cows. *J. Dairy Sci.*, 96,  
1384 6506-6513.  
1385
- 1386 Manninen, E., de Passillé, A. M., Rushen, J., Norring, M., and Saloniemi, H. 2002. Preferences  
1387 of dairy cows kept in unheated buildings for different kind of cubicle flooring. *Appl. Anim.*  
1388 *Behav. Sci.*, 75, 281-292.  
1389
- 1390 Manteuffel, G., Puppe, B., and Schön, P. C. 2004. Vocalization of farm animals as a measure of  
1391 welfare. *Appl. Anim. Behav. Sci.*, 88, 163-182.  
1392
- 1393 Manteuffel, G., Langbein, J., and Puppe, B. 2009. From operant learning to cognitive enrichment  
1394 in farm animal housing: bases and applicability. *Anim. Welf.*, 18, 87-95.  
1395



- 1396 March, M. D., Haskell, M. J., Chagunda, M. G. G., Langford, F. M., and Roberts, D. J. 2014.  
 1397 Current trends in British dairy management regimens. *J. Dairy Sci.*, 97, 7985-7994.  
 1398
- 1399 Marchant-Forde, J. N., Marchant-Forde, R. M., and Weary, D. M. 2002. Responses of dairy  
 1400 cows and calves to each other's vocalisations after early separation. *Appl. Anim. Behav. Sci.*, 78,  
 1401 19-28.  
 1402
- 1403 Mason, G. J., and Burn, C. C. 2011. Behavioural restriction. *Animal Welfare* (2nd), CAB  
 1404 International, Oxford, UK, 98-119.  
 1405
- 1406 McCowan, B., DiLorenzo, A. M., Abichandani, S., Borelli, C., and Cullor, J. S. 2002.  
 1407 Bioacoustic tools for enhancing animal management and productivity: effects of recorded calf  
 1408 vocalizations on milk production in dairy cows. *Appl. Anim. Behav. Sci.*, 77, 13-20.  
 1409
- 1410 McFarland, D. 1999. *Animal Behaviour* (3rd), Addison Wesley Longman, Reading, UK.  
 1411
- 1412 Meehan, C. L., and Mench, J. A. 2007. The challenge of challenge: can problem solving  
 1413 opportunities enhance animal welfare?. *Appl. Anim. Behav. Sci.*, 102, 246-261.  
 1414
- 1415 Mendl, M., Burman, O. H., and Paul, E. S. 2010. An integrative and functional frame work for  
 1416 the study of animal emotion and mood. *Proc. R. Soc. B.*, 277, 2895-2904.  
 1417

- 1418 Mitchell, K. D., Stookey, J. M., Lturnas, D. K., Watts, J. M., Haley, D. B., and Huyde, T. 2004.  
 1419 The effects of blindfolding on behavior and heart rate in beef cattle during restraint. *Appl. Anim.*  
 1420 *Behav. Sci.*, 85, 233-245.  
 1421
- 1422 Miller, K., and Wood-Gush, D. G. M. 1991. Some effects of housing on the social behaviour of  
 1423 dairy cows. *Anim. Prod.*, 53, 271-278.  
 1424
- 1425 Mintline, E. M., Stewart, M., Rogers, A. R., Cox, N. R., Verkerk, G. A., Stookey, J. M.,  
 1426 Webster, J.R. and Tucker, C. B. 2013. Play behavior as an indicator of animal welfare:  
 1427 Disbudding in dairy calves. *Appl. Anim. Behav. Sci.*, 144, 22-30.  
 1428
- 1429 Morgan, K. N., and Tromborg, C. T. 2007. Sources of stress in captivity. *Appl. Anim. Behav.*  
 1430 *Sci.*, 102, 262-302.  
 1431
- 1432 Mounier, L., Veissier, I., Andanson, S., Delval, E., and Boissy, A. 2006. Mixing at the beginning  
 1433 of fattening moderates social buffering in beef bulls. *Appl. Anim. Behav. Sci.*, 96, 185-200.  
 1434
- 1435 NAHMS. 2010. Dairy 2007. Facility Characteristics and Cow Comfort on U.S. Dairy  
 1436 Operations, 2007. Vol. #524.1210. USDA:APHIS:VS:CEAH, Fort Collins, CO.  
 1437 Accessed Oct. 3, 2015.  
 1438 [http://www.aphis.usda.gov/animal\\_health/nahms/dairy/downloads/dairy07/Dairy07\\_ir\\_F](http://www.aphis.usda.gov/animal_health/nahms/dairy/downloads/dairy07/Dairy07_ir_F)  
 1439 [acilities.pdf](http://www.aphis.usda.gov/animal_health/nahms/dairy/downloads/dairy07/Dairy07_ir_F)  
 1440

- 1441 Neisen, G., Wechsler, B., and Gygax, L. 2009. Effects of the introduction of single heifers or  
1442 pairs of heifers into dairy-cow herds on the temporal and spatial associations of heifers and cows.  
1443 *Appl. Anim. Behav. Sci.*, 119, 127-136.  
1444
- 1445 Newberry, R.C. 1993. The space-time continuum, and its relevance to farm animals. *Etologia*, 3,  
1446 219-234.  
1447
- 1448 Newberry, R. C. 1995. Environmental enrichment: increasing the biological relevance of captive  
1449 environments. *Appl. Anim. Behav. Sci.*, 44, 229-243.  
1450
- 1451 Newberry, R. C., and Swanson, J. C. 2008. Implications of breaking mother–young social bonds.  
1452 *Appl. Anim. Behav. Sci.*, 110, 3-23.  
1453
- 1454 Newby, N. C., Duffield, T. F., Pearl, D. L., Leslie, K. E., LeBlanc, S. J., and von Keyserlingk,  
1455 M. A. 2013. Use of a mechanical brush by Holstein dairy cattle around parturition. *J. Dairy Sci.*,  
1456 96, 2339-2344.  
1457
- 1458 Nicol, C. J., Caplen, G., Edgar, J., and Browne, W. J. 2009. Associations between welfare  
1459 indicators and environmental choice in laying hens. *Anim. Behav.*, 78, 413-424.  
1460
- 1461 Ninomiya, S. 2014. Satisfaction of farm animal behavioral needs in behaviorally restricted  
1462 systems: Reducing stressors and environmental enrichment. *Anim. Sci. J.* 85, 634-638.  
1463

- 1464 Ninomiya, S., and Sato, S. 2009. Effects of ‘Five freedoms’ environmental enrichment on the  
1465 welfare of calves reared indoors. *Anim. Sci.*, 80, 347-351.  
1466
- 1467 Ninomiya, S., and Sato, S. 2011. The assessment of the effect of presenting a companion's face  
1468 picture on social isolation stress using saliva sampling in cows. *Anim. Sci. J.*, 82, 787-791.  
1469
- 1470 Norring, M., Manninen, E., de Passillé, A. M., Rushen, J., and Saloniemi, H. 2010. Preferences  
1471 of dairy cows for three stall surface materials with small amounts of bedding. *J. Dairy Sci.*, 93,  
1472 70-74.  
1473
- 1474 Norring, M., Manninen, E., de Passillé, A. M., Rushen, J., Munksgaard, L., and Saloniemi, H.  
1475 2008. Effects of sand and straw bedding on the lying behavior, cleanliness, and hoof and hock  
1476 injuries of dairy cows. *J. Dairy Sci.*, 91, 570-576.  
1477
- 1478 Nitrates Action Programme and Phosphorus Regulations 2015-2018. Accessed Sept. 15, 2015.  
1479 [http://www.doeni.gov.uk/niea/water-home/agri\\_regs/nitrates-and-phosphorus-regulations-2015-](http://www.doeni.gov.uk/niea/water-home/agri_regs/nitrates-and-phosphorus-regulations-2015-)  
1480 2018.htm.  
1481
- 1482 O’Connell, N. E., Wicks, H. C., Carson, A. F., and McCoy, M. A. 2008. Influence of post-  
1483 calving regrouping strategy on welfare and performance parameters in dairy heifers. *Applied*  
1484 *Anim. Behav.Sci.*, 114, 319-329.  
1485

- 1486 Oltenacu, P.A., Algers, B., 2005. Selection for increased production and the welfare of dairy  
1487 cows: are new breeding goals needed? *Ambio* 34, 311-315.  
1488
- 1489 Pajor, E. A., Rushen, J., and de Passillé, A. M. B. 2003. Dairy cattle's choice of handling  
1490 treatments in a Y-maze. *Appl. Anim. Behav. Sci.*, 80, 93-107.  
1491
- 1492 Piller, C. A., Stookey, J. M., and Watts, J. M. 1999. Effects of mirror-image exposure on heart  
1493 rate and movement of isolated heifers. *Appl. Anim. Behav. Sci.*, 63, 93-102.  
1494
- 1495 Pilz, M., Fischer-Tenhagen, C., Thiele, G., Tinge, H., Lotz, F., and Heuwieser, W. 2012.  
1496 Behavioural reactions before and during vaginal examination in dairy cows. *Appl. Anim. Behav.*  
1497 *Sci.*, 138, 18-27.  
1498
- 1499 Pollock, W. E., and Hurnik, J. F. 1978. Effect of calf calls on rate of milk release of dairy cows.  
1500 *J. Dairy Sci.*, 61, 1624-1626.  
1501
- 1502 Popescu, S., Borda, C., Diugan, E. A., Spinu, M., Groza, I. S., and Sandru, C. D. 2013. Dairy  
1503 cows welfare quality in tie-stall housing system with or without access to exercise. *Acta Vet.*  
1504 *Scand.*, 55, 43.  
1505
- 1506 Probst, J. K., Neff, A. S., Leiber, F., Kreuzer, M., and Hillmann, E. 2012. Gentle touching in  
1507 early life reduces avoidance distance and slaughter stress in beef cattle. *Appl. Anim. Behav. Sci.*,  
1508 139, 42-49.

1509

1510 Proctor, H.S., and Carder, G. 2014. Can ear postures reliably measure the positive emotional

1511 state of cows?. *Appl. Anim. Behav. Sci.*, 161, 20-27.

1512

1513 Proctor, H.S, and Carder, G. 2015a. Measuring positive emotions in cows: Do visible eye whites

1514 tell us anything? *Physiol. Behav.*, 147, 1-6.

1515

1516 Proctor, H. S., and Carder, G. 2015b. Nasal temperatures in dairy cows are influenced by

1517 positive emotional state. *Physiol. Behav.*, 138, 340-344.

1518

1519 Proudfoot, K. L., Jensen, M. B., Weary, D. M., and von Keyserlingk, M. A. G. 2014b. Dairy

1520 cows seek isolation at calving and when ill. *J. Dairy Sci.*, 97, 2731-2739.

1521

1522 Proudfoot, K. L., Weary, D. M., and von Keyserlingk, M. A. G. 2014a. Maternal isolation

1523 behavior of Holstein dairy cows kept indoors. *Anim. Sci. J.*, 92, 277-281.

1524

1525 Rault, J. L. 2012. Friends with benefits: Social support and its relevance for farm animal welfare.

1526 *Appl. Anim. Behav. Sci.*, 136, 1-14.

1527

1528 Raussi, S., Niskanen, S., Siivonen, J., Hänninen, L., Hepola, H., Jauhiainen, L., and Veissier, I.

1529 2010. The formation of preferential relationships at early age in cattle. *Behav. Processes*, 84,

1530 726-731.

1531

- 1532 Rauw, W. M. et al., 1998. Undesirable side effects of selection for high production efficiency in  
1533 farm animals: a review. *Livest. Prod. Sci.* 56, 15-33  
1534
- 1535 Reijs, J. W., C. H. G. Daatselaar, J. F. M. Helming, J. Jager, and A. C. G. Beldman. 2013.  
1536 Grazing dairy cows in north-west Europe. Economic farm performance and future developments  
1537 with emphasis on the Dutch situation. Report of the Landbouw Economisch Instituut (LEI). LEI  
1538 Wageningen UR, Wageningen, the Netherlands.  
1539
- 1540 Regula, G., Danuser, J., Spycher, B., and Wechsler, B. 2004. Health and welfare of dairy cows in  
1541 different husbandry systems in Switzerland. *Prev. Vet. Med.* 66, 247-264.  
1542
- 1543 Rioja-Lang, F. C., Roberts, D. J., Healy, S. D., Lawrence, A. B., and Haskell, M. J. 2009. Dairy  
1544 cows trade-off feed quality with proximity to a dominant individual in Y-maze choice tests.  
1545 *Appl. Anim. Behav. Sci.*, 117, 159-164.  
1546
- 1547 Roth, B. A., Barth, K., Gygax, L., and Hillmann, E. 2009. Influence of artificial vs. mother-  
1548 bonded rearing on sucking behaviour, health and weight gain in calves. *Appl. Anim. Behav. Sci.*,  
1549 119, 143-150.  
1550
- 1551 Rushen, J., Munksgaard, L., de Passillé, A. M. B., Jensen, M. B., and Thodberg, K. 1998.  
1552 Location of handling and dairy cows' responses to people. *Appl. Anim. Behav. Sci.*, 55, 259-267.  
1553

1554 Rushen, J., Munksgaard, L., Marnet, P. G., and de Passillé, A. M. 2001. Human contact and the  
1555 effects of acute stress on cows at milking. *Appl. Anim. Behav. Sci.*, 73, 1-14.

1556  
1557 Rushen, J., Taylor, A. A., and de Passillé, A. M. 1999. Domestic animals' fear of humans and its  
1558 effect on their welfare. *Appl. Anim. Behav. Sci.*, 65, 285-303.

1559  
1560 Sambrook, T. D., and Buchanan-Smith, H. M. 1997. Control and complexity in novel object  
1561 enrichment. *Anim. Welf.*, 6, 207-216.

1562  
1563 Sandem, A. I., Braastad, B. O., and Bøe, K. E. 2002. Eye white may indicate emotional state on a  
1564 frustration–contentedness axis in dairy cows. *Appl. Anim. Behav. Sci.*, 79, 1-10.

1565  
1566 Sandem, A. I., Janczak, A. M., and Braastad, B. O. 2004. A short note on effects of exposure to a  
1567 novel stimulus (umbrella) on behaviour and percentage of eye-white in cows. *Appl. Anim.*  
1568 *Behav. Sci.*, 89, 309-314.

1569  
1570 Sato, S., Sako, S., and Maeda, A. 1991. Social licking patterns in cattle (*Bos taurus*): influence of  
1571 environmental and social factors. *Appl. Anim. Behav. Sci.*, 32, 3-12.

1572  
1573 Sato, S., Tarumizu, K., and Hatae, K. 1993. The influence of social factors on allogrooming in  
1574 cows. *Appl. Anim. Behav. Sci.*, 38, 235-244.

1575



- 1576 Schirmann, K., Chapinal, N., Weary, D. M., Heuwieser, W., and von Keyserlingk, M. A. G.  
 1577 2011. Short-term effects of regrouping on behavior of prepartum dairy cows. *J. Dairy Sci.*, 94,  
 1578 2312-2319.
- 1579
- 1580 Schloeth, R. 1961. Das socialleben des carmargue Rindes. *Z. Tierpsychol.*, 18, 574–627.
- 1581
- 1582 Schukken, Y. H., and Young, G. D. 2009. Field study on milk production and mastitis effect of  
 1583 the DeLaval Swinging Cow Brush. DeLaval Swinging Cow Brush Study Final Report, August 5.  
 1584 DeLaval, Tumba, Sweden.
- 1585
- 1586 Schütz, K. E., Clark, K. V., Cox, N. R., Matthews, L. R., and Tucker, C. B. 2010. Responses to  
 1587 short-term exposure to simulated rain and wind by dairy cattle: time budgets, shelter use, body  
 1588 temperature and feed intake. *Anim. Welf.*, 19, 375.
- 1589
- 1590 Schütz, K. E., Hawke, M., Waas, J. R., McLeay, L. M., Bokkers, E. A. M., van Reenen, C. G.,  
 1591 Webster, J.R and Stewart, M. 2012. Effects of human handling during early rearing on the  
 1592 behaviour of dairy calves. *Anim. Welf.*, 21, 19-26.
- 1593
- 1594 Shyne, A. 2006. Meta-analytic review of the effects of enrichment on stereotypic behavior in zoo  
 1595 mammals. *Zoo Biol.*, 25, 317-337.
- 1596
- 1597 Spinka, M., and Wemelsfelder, F. 2011. Environmental challenge and animal agency. *Animal*  
 1598 *Welfare* (2nd). CABI International, Wallingford, 27-43.

- 1599
- 1600 Stanford, K., Croy, D., Bach, S. J., Wallins, G. L., Zahiroddini, H., and McAllister, T. A. 2005.
- 1601 Ecology of *Escherichia coli* O157: H7 in Commercial Dairies in Southern Alberta. J. Dairy Sci.,
- 1602 88, 4441-4451.
- 1603
- 1604 Stanford, K., Silasi, R., McAllister, T. A., and Schwartzkopf-Genswein, K. S. 2009. Behavior of
- 1605 feedlot cattle affects voluntary oral and physical interactions with manila ropes. Anim. Sci. J.,
- 1606 87, 296-303.
- 1607
- 1608 Stewart, M., Shepherd, H. M., Webster, J. R., Waas, J. R., McLeay, L. M., and Schütz, K. E.
- 1609 2013. Effect of previous handling experiences on responses of dairy calves to routine husbandry
- 1610 procedures. Animal, 7, 828-833.
- 1611
- 1612 Svensson, C., and Liberg, P. 2006. The effect of group size on health and growth rate of Swedish
- 1613 dairy calves housed in pens with automatic milk-feeders. Prev. Vet. Med., 73, 43-53.
- 1614
- 1615 Svensson, C., Hultgren, J., and Oltenacu, P. A. 2006. Morbidity in 3–7-month-old dairy calves in
- 1616 south-western Sweden, and risk factors for diarrhoea and respiratory disease. Prev. Vet. Med.,
- 1617 74, 162-179.
- 1618
- 1619 Svensson, C., Lundborg, K., Emanuelson, U., and Olsson, S. O. 2003. Morbidity in Swedish
- 1620 dairy calves from birth to 90 days of age and individual calf-level risk factors for infectious
- 1621 diseases. Prev. Vet. Med., 58, 179-197.

1622

1623 Talebi, A., von Keyserlingk, M. A. G., Telezhenko, E., and Weary, D. M. 2014. Reduced  
1624 stocking density mitigates the negative effects of regrouping in dairy cattle. *J. Dairy Sci.*, 97,  
1625 1358-1363.

1626

1627 Taylor, A. A., and Davis, H. 1998. Individual humans as discriminative stimuli for cattle (*Bos*  
1628 *taurus*). *Appl. Anim. Behav. Sci.*, 58, 13-21.

1629

1630 Terlouw, E. M., Boissy, A., and Blinet, P. 1998. Behavioural responses of cattle to the odours of  
1631 blood and urine from conspecifics and to the odour of faeces from carnivores. *Appl. Anim.*  
1632 *Behav. Sci.*, 57, 9-21.

1633

1634 Thomas, T. J., Weary, D. M., and Appleby, M. C. 2001. Newborn and 5-week-old calves  
1635 vocalize in response to milk deprivation. *Appl. Anim. Behav. Sci.*, 74, 165-173.

1636

1637 Toaff-Rosenstein, R. L., Gershwin, L. J., Zanella, A. J., Tucker, C. B. 2014. Characterizing the  
1638 BRD sickness response - opportunities for improved disease detection. 4th International  
1639 symposium on beef cattle welfare, Ames, Iowa.

1640

1641 Tucker, C. B. 2009. Behaviour of cattle. pp 1151. Jensen, P. (Ed.). *The Ethology of Domestic*  
1642 *Animals: an introductory text*. CAB International, Wallingford.

1643

- 1644 Tucker, C. B., Weary, D. M., and Fraser, D. 2003. Effects of three types of free-stall surfaces on  
1645 preferences and stall usage by dairy cows. *J. Dairy Sci.*, 86, 521-529.  
1646
- 1647 Tuytens, F. A. M. 2005. The importance of straw for pig and cattle welfare: a review. *Appl.*  
1648 *Anim. Behav. Sci.*, 92, 261-282  
1649
- 1650 Ude, G., Georg, H., and Schwalm, A. 2011. Reducing milk induced cross-sucking of group  
1651 housed calves by an environmentally enriched post feeding area. *Livest. Sci.*, 138, 293-298.  
1652
- 1653 Uetake, K., Hurnik, J. F., and Johnson, L. 1997. Effect of music on voluntary approach of dairy  
1654 cows to an automatic milking system. *Appl. Anim. Behav. Sci.*, 53, 175-182.  
1655
- 1656 Val-Laillet, D., Veira, D. M., and von Keyserlingk, M. A. G. 2008. Dominance in Free-Stall—  
1657 Housed Dairy Cattle Is Dependent upon Resource. *J. Dairy Sci.*, 91, 3922-3926.  
1658
- 1659 Val-Laillet, D., Guesdon, V., von Keyserlingk, M. A., de Passillé, A. M., and Rushen, J. 2009.  
1660 Allogrooming in cattle: relationships between social preferences, feeding displacements and  
1661 social dominance. *Appl. Anim. Behav. Sci.*, 116, 141-149.  
1662
- 1663 Valníčková, B., Stěhulová, I., Šárová, R., and Špinka, M. 2015. The effect of age at separation  
1664 from the dam and presence of social companions on play behavior and weight gain in dairy  
1665 calves. *J. Dairy Sci.*, 98, 5545-5556.  
1666

- 1667 van de Weerd, H. A., and Day, J. E. 2009. A review of environmental enrichment for pigs  
1668 housed in intensive housing systems. *Appl. Anim. Behav. Sci.*, 116, 1-20.  
1669
- 1670 Van Vuuren, A. M., and Van den Pol-van Dasselaar, A. 2007. Grazing systems and feed  
1671 supplementation. *Frontis*, 18, 85-101.  
1672
- 1673 Vaughan, A., de Passillé, A. M., Stookey, J., and Rushen, J. 2014. Operant conditioning of  
1674 urination by calves. *Appl. Anim. Behav. Sci.*, 158, 8-15.  
1675
- 1676 Veerkamp, R.F. et al., 2009. Selection for high production in dairy cattle. In: Rauw, W.M. (Ed.),  
1677 Resource Allocation Theory Applied to Farm Animal Production. CAB International,  
1678 Wallingford, Oxfordshire, UK, pp. 243-260.  
1679
- 1680 Veissier, I., Andanson, S., Dubroeuq, H., and Pomiès, D. 2008. The motivation of cows to walk  
1681 as thwarted by tethering. *Anim. Sci. J.*, 86, 2723-2729.  
1682
- 1683 Ventura, B. A., Von Keyserlingk, M. A. G., Schuppli, C. A., and Weary, D. M. 2013. Views on  
1684 contentious practices in dairy farming: The case of early cow-calf separation. *J. Dairy. Sci.*, 96,  
1685 6105-6116.  
1686
- 1687 von Keyserlingk, M. A. G., Olenick, D., and Weary, D. M. 2008. Acute behavioral effects of  
1688 regrouping dairy cows. *J. Dairy Sci.*, 91, 1011-1016.  
1689

- 1690 Wagner, K., Barth, K., Palme, R., Futschik, A., and Waiblinger, S. 2012. Integration into the  
1691 dairy cow herd: long-term effects of mother contact during rearing the first twelve weeks of life.  
1692 Appl. Anim. Behav. Sci. 141, 117-129.  
1693
- 1694 Wagner, K., Barth, K., Hillmann, E., Palme, R., Futschik, A., and Waiblinger, S. 2013. Mother  
1695 rearing of dairy calves: reactions to isolation and to confrontation with an unfamiliar conspecific  
1696 in a new environment. Appl. Anim. Behav. Sci. 147, 43-54.  
1697
- 1698 Wagner, K., Seitner, D., Barth, K., Palme, R., Futschik, A., and Waiblinger, S. 2015. Effects of  
1699 mother versus artificial rearing during the first 12 weeks of life on challenge responses of dairy  
1700 cows. Appl. Anim. Behav. Sci. 164, 1-11.  
1701
- 1702 Waiblinger, S., Boivin, X., Pedersen, V., Tosi, M. V., Janczak, A. M., Visser, E. K., and Jones,  
1703 R. B. 2006. Assessing the human–animal relationship in farmed species: a critical review. Appl.  
1704 Anim. Behav. Sci., 101, 185-242.  
1705
- 1706 Waiblinger, S., Menke, C., and Coleman, G. 2002. The relationship between attitudes, personal  
1707 characteristics and behaviour of stockpeople and subsequent behaviour and production of dairy  
1708 cows. Appl. Anim. Behav. Sci., 79, 195-219.  
1709
- 1710 Waiblinger, S., Menke, C., Korff, J., and Bucher, A. 2004. Previous handling and gentle  
1711 interactions affect behaviour and heart rate of dairy cows during a veterinary procedure. Appl.  
1712 Anim. Behav. Sci., 85, 31-42.

- 1713
- 1714 Warnick, V. D., Arave, C. W., and Mickelsen, C. H. 1977. Effects of group, individual, and
- 1715 isolated rearing of calves on weight gain and behavior. *J. Dairy Sci.*, 60, 947-953.
- 1716
- 1717 Waynert, D. F., Stookey, J. M., Schwartzkopf-Genswein, K. S., Watts, J. M., and Waltz, C. S.
- 1718 1999. The response of beef cattle to noise during handling. *Appl. Anim. Behav. Sci.*, 62, 27-42.
- 1719
- 1720 Weary, D. M., and Chua, B. 2000. Effects of early separation on the dairy cow and calf.: 1.
- 1721 Separation at 6 h, 1 day and 4 days after birth. *Appl. Anim. Behav. Sci.*, 69, 177-188.
- 1722
- 1723 Webb, L.E., Jensen, M.B., Engel, B., van Reenen, C.G., Gerrits, W.J.J., de Boer, I.J.M., Bokkers,
- 1724 E.A.M., 2014b. Chopped or long roughage: what do calves prefer? Using cross point analysis of
- 1725 double demand functions. *PLoS One* 9, e88778.
- 1726
- 1727 Wechsler, B., and Lea, S. E. 2007. Adaptation by learning: its significance for farm animal
- 1728 husbandry. *Appl. Anim. Behav. Sci.*, 108, 197-214.
- 1729
- 1730 Wells, D. L. 2009. Sensory stimulation as environmental enrichment for captive animals: a
- 1731 review. *Appl. Anim. Behav. Sci.*, 118, 1-11.
- 1732
- 1733 Wells, S. J., Garber, L. P., and Hill, G. W. 1997. Health status of preweaned dairy heifers in the
- 1734 United States. *Prev. Vet. Med.*, 29, 185-199.
- 1735

1736 Westerath, H.S., Gyax, L., Hillmann, E. 2014. Are special feed and being brushed judged as  
1737 positive by calves?, *Appl. Anim. Behav. Sci.*, 156, 12-21.

1738

1739 Wilson, S. C., Mitlöhner, F. M., Morrow-Tesch, J., Dailey, J. W., and McGlone, J. J. 2002. An  
1740 assessment of several potential enrichment devices for feedlot cattle. *Appl. Anim. Behav. Sci.*,  
1741 76, 259-265.

1742

1743 Winsten, J. R., Kerchner, C. D., Richardson, A., Lichau, A., and Hyman, J. M. 2010. Trends in  
1744 the Northeast dairy industry: Large-scale modern confinement feeding and management-  
1745 intensive grazing. *J. Dairy Sci.*, 93, 1759-1769.

1746

1747 Wood-Gush, D. G. M., Hunt, K., Carson, K., and Dennison, S. G. C. 1984. The early behaviour  
1748 of suckler calves in the field. *Biol. Behav.*, 9, 295-306.

1749

1750 Wredle, E., Munksgaard, L., and Spöndly, E. 2006. Training cows to approach the milking unit  
1751 in response to acoustic signals in an automatic milking system during the grazing season. *Appl.*  
1752 *Anim. Behav. Sci.*, 101, 27-39.

1753

1754 Wredle, E., Rushen, J., de Passille, A. M., and Munksgaard, L. 2004. Training cattle to approach  
1755 a feed source in response to auditory signals. *Can. J. Anim. Sci.*, 84, 567-572.